

HIF1-alpha ABfinity™ Recombinant Rabbit Monoclonal Antibody - Purified



Catalog no. 700505

(See product label for lot information)

Clone/PAD: 16H4L13
Isotype: IgG
Gene ID: 3091
Protein Acc. no.: Q16665
Qty: 100 µg
Volume: 200 µl
Concentration: 0.5 mg/mL

Formulation

Phosphate buffered saline (PBS) with 0.09% sodium azide.

Application

For use in Western Blotting and ELISA.

Reactivity

This antibody is specific for human HIF1-alpha.

Immunogen

Peptide

Immunogen sequence

EDTEAKNPFSTQ

Sequence Identity

Human

Sequence Homology

Mouse

Expected Reactivity

Based on sequence identity and similarity, reactivity to Human and Mouse are predicted.

Storage

2-8°C for up to 1 month, -20°C for long term storage. Avoid repeated freezing and thawing.

Expiration Date

Expires one year from date of receipt when stored as instructed.

Background

HIF1-alpha (Hypoxia-inducible factor 1, alpha), a subunit HIF1, which is a transcription factor found in mammalian cells cultured under reduced oxygen tension (1). HIF1 functions as a transcriptional regulator of the adaptive response to hypoxia. HIF1-alpha regulates hypoxia-mediated apoptosis, cell proliferation and tumour angiogenesis (2). Hypoxia which induces p53 protein accumulation, directly interacts with HIF1-alpha and reduces hypoxia-induced expression of HIF1-alpha by promoting MDM2-mediated ubiquitination and proteasomal degradation under hypoxic conditions (3-4). Recent studies suggests that induction of NOX4 by HIF1-alpha contributes to maintain ROS levels after hypoxia and hypoxia-induced proliferation (5). In humans, it is located on the q arm of chromosome 14 (6).

References

1. Wang GL, Jiang BH, Rue EA, Semenza GL. (1995). Hypoxia-inducible factor 1 is a basic-helix-loop-helix-PAS heterodimer regulated by cellular O2 tension. *Proceeding of the National Academy of Science U S A.*, 92(12):5510-4.
2. Carmeliet P, Dor Y, Herbert JM, Fukumura D, Brusselmans K, Dewerchin M, Neeman M, Bono F, Abramovitch R, Maxwell P, Koch CJ, Ratcliffe P, Moons L, Jain RK, Collen D, Keshert E. (1998). Role of HIF-1alpha in hypoxia-mediated apoptosis, cell proliferation and tumour angiogenesis. *Nature*, 394(6692):485-90.
3. Chen D, Li M, Luo J, Gu W. (2003). Direct interactions between HIF-1 alpha and Mdm2 modulate p53 function. *The Journal of Biological chemistry*, 278(16):13595-8.
4. Anna-Liisa Nieminen, Suparna Qanungo, Elizabeth A. Schneider, Bing-Hua Jiang, Faton H. Agani. (2005). Mdm2 and HIF-1 interaction in tumor cells during hypoxia. *Journal of Cellular Physiology*, 364-369.
5. Diebold I, Petry A, Hess J, Görlach A. (2010). The NADPH oxidase subunit NOX4 is a new target gene of the hypoxia-inducible factor-1. *Molecular Biology of the Cell*, 21(12):2087-96.
6. Semenza GL, Rue EA, Iyer NV, Pang MG, Kearns WG. (1996). Assignment of the hypoxia-inducible factor 1alpha gene to a region of conserved syntenry on mouse chromosome 12 and human chromosome 14q. *Genomics*, 34(3):437-9.



Applications:

	Species	Test Material	Concentration
Western Blotting	Human	HEK	0.1 - 2 µg/ml
Indirect ELISA	Human	HEK	5x10 ⁻⁴ - 10 µg/ml

For Research Use Only. CAUTION: Not for human or animal therapeutic or diagnostic use.

www.invitrogen.com

Invitrogen Corporation • 7335 Executive Way • Frederick • MD 21704 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com



Western Blot analysis of HIF1-alpha ABfinity™ Recombinant Rabbit Monoclonal Antibody (Cat. No.700505).

Whole cell extract; 30µg per lane from HEK was loaded on SDS-PAGE followed by transfer on to nitrocellulose. The blot was blocked followed by incubation with HIF1-alpha ABfinity™ Recombinant Rabbit Monoclonal Antibody at 0.1µg/mL for 2 hours. Goat Anti Rabbit - HRP conjugated, was used at 1:5000 dilution as secondary antibody and developed by chemiluminescence (ECL) method. Expected size is ~93kDa.

Indirect ELISA of Hif1-alpha Rabbit Recombinant Monoclonal Antibody (Cat. No.700505).

Indirect ELISA was done using Hif1 alpha Rabbit Recombinant Monoclonal Antibody to detect Hif1 alpha Rabbit Recombinant Monoclonal Antibody on HEK293 cell lysate (1ug/well) using TMB (Cat. No. SB01) as substrate.

Explanation of symbols

Symbol	Description	Symbol	Description
	Catalogue Number		Batch code
	Research Use Only		In vitro diagnostic medical device
	Use by		Temperature limitation
	Manufacturer		European Community authorised representative
	Without, does not contain		With, contains
	Protect from light		Consult accompanying documents
	Directs the user to consult instructions for use (IFU), accompanying the product.		

Limited Use Label License No. 358: Research Use Only

The purchase of this product conveys to the purchaser the limited, non-transferable right to use the purchased amount of the product only to perform internal research for the sole benefit of the purchaser. No right to resell this product or any of its components is conveyed expressly, by implication, or by estoppel. This product is for internal research purposes only and is not for use in commercial applications of any kind, including, without limitation, quality control and commercial services such as reporting the results of purchaser's activities for a fee or other form of consideration. For information on obtaining additional rights, please contact outlicensing@lifetech.com or Out Licensing, Life Technologies, 5791 Van Allen Way, Carlsbad, California 92008.

©2011 Life Technologies Corporation. All rights reserved. The trademarks mentioned herein are the property of Life Technologies Corporation or their respective owners.

For Research Use Only. CAUTION: Not for human or animal therapeutic or diagnostic use.

www.invitrogen.com

Invitrogen Corporation • 7335 Executive Way • Frederick • MD 21704 • Tel: 800.955.6288 • E-mail: techsupport@invitrogen.com