

Contents

1. Description
 - 1.1 Background information
 - 1.2 Applications
2. References

1. Description

Products	Human IL-12, premium grade. Recombinant human interleukin 12.								
	<table border="1"> <thead> <tr> <th>Content in µg</th> <th>Order no.</th> </tr> </thead> <tbody> <tr> <td>5</td> <td>130-096-704</td> </tr> <tr> <td>25</td> <td>130-096-705</td> </tr> <tr> <td>100</td> <td>130-096-798</td> </tr> </tbody> </table>	Content in µg	Order no.	5	130-096-704	25	130-096-705	100	130-096-798
Content in µg	Order no.								
5	130-096-704								
25	130-096-705								
100	130-096-798								
Biological activity	The ED ₅₀ is ≤0.33 ng/mL corresponding to an activity of ≥3×10 ⁶ U/mg. For lot-specific activities, please contact our technical support. ▲ Note: The ED ₅₀ is determined by inducing of IFN-γ secretion by PHA-activated T cells. ¹ The proliferation assay was calibrated with the reference reagent for human IL-12 (NIBSC code 95/544) provided by the WHO/National Institute for Biological Standards and Control.								
Primary structure	Glycosylated single-chain polypeptide, p35 and p40 fused by a flexible linker region (531 amino acid residues).								
Molecular mass	60 kDa (calculated). 80 kDa (SDS-PAGE under reducing conditions).								
Source	Produced in HEK293 cells.								
Product format	Lyophilized from a filtered (0.2 µm) buffer solution.								
Stabilizer	Mannitol and trehalose.								
Purity	>97% as determined by SDS-PAGE analysis.								
Endotoxin level	Low endotoxin (<0.1 EU/µg cytokine) as determined by Limulus Amebocyte Lysate (LAL) assay.								
Storage	Lyophilized Human IL-12, premium grade should be stored at -20 °C. The expiration date is indicated on the vial label. Upon reconstitution aliquots should be stored at -20 °C or below. Avoid repeated freeze-thaw cycles.								
Reconstitution	It is recommended to reconstitute lyophilized Human IL-12, premium grade with deionized sterile-filtered water to a final concentration of 0.05–1.0 mg/mL in a minimal volume of 100 µL. Further dilutions should be prepared with 0.1% bovine serum albumin (BSA) or human serum albumin (HSA) in phosphate-buffered saline.								

1.1 Background information

IL-12 is a heterodimeric proinflammatory cytokine and a modulator of cell-mediated immunity, which is mainly produced by macrophages, dendritic cells, and B cells. IL-12 stimulates the production and secretion of several cytokines, in particular IFN-γ, by NK cells and T cells, induces proliferation, and enhances the cytotoxic activity within these cell populations. Another important activity of IL-12, acting together with IFN-γ and IL-2, is to drive T helper cell responses toward the T_H1 rather than the T_H2 phenotype. Moreover, IL-12 is also important in resistance to viral disease and has significant antitumor activity. It has been shown that single chain fusion proteins of naturally occurring heterodimeric cytokines such as IL-12 or IL-23 are bioactive *in vitro* and *in vivo*.

1.2 Applications

Human IL-12 can be used for a variety of applications, including:

- *In vitro* differentiation of naive CD4⁺ T cells towards T_H1 cells.
- *In vitro* proliferation and stimulation of cytotoxic activity of NK cells and T cells.

Optimal concentration for a specific application should be determined by a dose-response experiment.

2. References

1. Wulff, H. *et al.* (2007) Cloning and characterization of an adenoviral vector for highly efficient and doxycycline-suppressible expression of bioactive human single-chain interleukin 12 in colon cancer. *BMC Biotechnol.* 7: 35.
2. Lieschke, G.J. *et al.* (1997) Bioactive murine and human interleukin-12 fusion proteins which retain antitumor activity *in vivo*. *Nat. Biotechnol.* 15: 35–40.
3. Miller, J.M. *et al.* (2010) Vesicular stomatitis virus modified with single chain IL-23 exhibits oncolytic activity against tumor cells *in vitro* and *in vivo*. *Int J Inference Cytokine Mediator Res* 2010: 63–72.

Refer to www.miltenyibiotec.com for all data sheets and protocols.

Warranty

The products sold hereunder are warranted only to be free from defects in workmanship and material at the time of delivery to the customer. Miltenyi Biotec GmbH makes no warranty or representation, either expressed or implied, with respect to the fitness of a product for a particular purpose. There are no warranties, expressed or implied, which extend beyond the technical specifications of the products. Miltenyi Biotec GmbH's liability is limited to either replacement of the products or refund of the purchase price. Miltenyi Biotec GmbH is not liable for any property damage, personal injury or economic loss caused by the product.

MACS is a registered trademark of Miltenyi Biotec GmbH.

Unless otherwise specifically indicated, Miltenyi Biotec products and services are for research use only and not for diagnostic or therapeutic use.

Copyright © 2015 Miltenyi Biotec GmbH. All rights reserved.