

Anti-TCR Vβ17 antibodies, human

For research use only

One test corresponds to labeling of up to 10⁶ cells in a total volume of 100 μL

| Product | Content | Order no. |
|--------------------------|---------------|-------------|
| Anti-TCR Vβ17-Biotin | for 100 tests | 130-115-246 |
| Anti-TCR Vβ17-FITC | for 30 tests | 130-115-346 |
| Anti-TCR Vβ17-FITC | for 100 tests | 130-115-247 |
| Anti-TCR Vβ17-PE | for 30 tests | 130-115-347 |
| Anti-TCR Vβ17-PE | for 100 tests | 130-115-248 |
| Anti-TCR Vβ17-APC | for 30 tests | 130-115-348 |
| Anti-TCR Vβ17-APC | for 100 tests | 130-115-249 |
| Anti-TCR Vβ17-PE-Vio770 | for 30 tests | 130-115-349 |
| Anti-TCR Vβ17-PE-Vio770 | for 100 tests | 130-115-250 |
| Anti-TCR Vβ17-APC-Vio770 | for 30 tests | 130-115-350 |
| Anti-TCR Vβ17-APC-Vio770 | for 100 tests | 130-115-251 |
| Anti-TCR Vβ17-Biotin | for 30 tests | 130-115-345 |

Warnings

Reagents contain sodium azide. Under acidic conditions sodium azide yields hydrazoic acid, which is extremely toxic. Azide compounds should be diluted with running water before discarding. These precautions are recommended to avoid deposits in plumbing where explosive conditions may develop.

Technical data and background information

| | |
|--|---|
| Antigen | TCR Vβ17 |
| Clone | REA915 |
| Isotype | recombinant human IgG1 |
| Isotype control | REA Control (S) antibodies |
| Alternative names of antigen | TCR Vbeta17, TCR Vb17 |
| Entrez Gene ID | 28570 |
| Molecular mass of antigen [kDa] | 11 |
| Distribution of antigen | T cells |
| Product format | Reagents are supplied in buffer containing stabilizer and 0.05% sodium azide. |
| Fixation | Cells should be stained prior to fixation, if formaldehyde is used as a fixative. |
| Storage | Store protected from light at 2-8 °C. Do not freeze. |

Clone REA915 recognizes the human variable β 17 subunit of the αβ T cell receptor (TCR Vβ17). The TCR is a heterodimeric glycoprotein associated with the CD3 antigen. The α and β TCR chains are composed of constant and variable regions, each encoded by distinct gene segments. TCR Vβ17 is a variant of the TCR β chain. αβ T cells express a sparse αβ TCR repertoire

that specifically co-recognizes self or foreign antigen bound to antigen-presenting molecules, which thereby leads to T cell-mediated immunity. For example, the TCR can directly bind to peptide fragments, riboflavin precursors, and lipid antigens that are presented by major histocompatibility complex (MHC) molecules, MR1 and CD1, respectively. In each case, the antigen sits within the antigen-binding cleft, whereupon the TCR recognizes a composite surface formed by the antigen-presenting molecule and surface-exposed regions of the antigen itself. This co-recognition paradigm is a central tenet of $\alpha\beta$ T cell-mediated immunity and underpins MHC restriction. Additional information: Clone REA915 displays negligible binding to Fc receptors.

Reagent requirements











- Buffer: Prepare a solution containing phosphate-buffered saline (PBS), pH 7.2, 0.5% bovine serum albumin (BSA), and 2 mM EDTA by diluting MACS[®] BSA Stock Solution (# 130-091-376) 1:20 with autoMACS[®] Rinsing Solution (# 130-091-222). Keep buffer cold (2–8 °C).
Note: EDTA can be replaced by other supplements such as anticoagulant citrate dextrose formula-A (ACD-A) or citrate phosphate dextrose (CPD). Buffers or media containing Ca²⁺ or Mg²⁺ are not recommended for use.
- (Optional) Fluorochrome-conjugated anti-biotin antibodies, e.g., Anti-Biotin-PE (# 130-090-756) as secondary antibody reagent in combination with biotinylated antibodies.
- (Optional) Propidium Iodide Solution (# 130-093-233) for flow cytometric exclusion of dead cells without fixation.
- (Optional) Fixation and Dead Cell Discrimination Kit (# 130-091-163) for cell fixation and flow cytometric exclusion of dead cells.



Protocol for cell surface staining

- The recommended antibody dilution for labeling of cells and subsequent analysis by flow cytometry is 1:50 for up to 10⁶ cells/100 μ L.
 - Volumes given below are for up to 10⁶ nucleated cells. When working with fewer than 10⁶ cells, use the same volumes as indicated. When working with higher cell numbers, scale up all reagent volumes and total volumes accordingly.
1. Determine cell number.
 2. Centrifuge cell suspension at 300 \times g for 10 minutes. Aspirate supernatant completely.
 3. Resuspend up to 10⁶ nucleated cells per 98 μ L of buffer.
 4. Add 2 μ L of the antibody.
 5. Mix well and incubate for 10 minutes in the dark in the refrigerator (2–8 °C).
Note: Higher temperatures and/or longer incubation times may lead to non-specific cell labeling. Working on ice requires increased incubation times.
 6. Wash cells by adding 1–2 mL of buffer and centrifuge at 300 \times g for 10 minutes. Aspirate supernatant completely.
 7. (Optional) If biotinylated antibody was used, resuspend the cell pellet in buffer and stain with fluorochrome-conjugated anti-biotin antibody according to the manufacturer's recommendations.
 8. Resuspend cell pellet in a suitable amount of buffer for analysis by flow cytometry or fluorescence microscopy.

Examples of immunofluorescent staining

Human peripheral blood mononuclear cells (PBMCs) were stained with Anti-TCR V β 17 antibodies or with the corresponding REA Control (S) antibodies (left images) as well as with CD3 antibodies. Flow cytometry was performed using the MACSQuant[®] Analyzer. The Tandem Signal Enhancer has been used to increase binding specificity of tandem-dye-conjugated antibodies. Cell debris and dead cells were excluded from the analysis based on scatter signals and propidium iodide fluorescence or 4',6-diamidino-2-phenylindole (DAPI) fluorescence, as in the case of tandem conjugates.





Warranty

The products sold hereunder are warranted only to be free from defects in workmanship and material at the time of delivery to the customer. Miltenyi Biotec GmbH makes no warranty or representation, either expressed or implied, with respect to the fitness of a product for a particular purpose. There are no warranties, expressed or implied, which extend beyond the technical specifications of the products. Miltenyi Biotec GmbH's liability is limited to either replacement of the products or refund of the purchase price. Miltenyi Biotec GmbH is not liable for any property damage, personal injury or economic loss caused by the product.

Miltenyi Biotec GmbH | Friedrich-Ebert-Straße 68 | 51429 Bergisch Gladbach | Germany | Phone +49 2204 8306-0 | Fax +49 2204 85197 | macs@miltenyibiotec.de | www.miltenyibiotec.com Miltenyi Biotec provides products and services worldwide. Visit www.miltenyibiotec.com/local to find your nearest Miltenyi Biotec contact.

Unless otherwise specifically indicated, Miltenyi Biotec products and services are for research use only and not for therapeutic or diagnostic use. autoMACS, MACS, MACSQuant, Vio, VioBlue, VioBright, and VioGreen are either trademarks or registered trademarks of Miltenyi Biotec GmbH. Copyright © 2016 Miltenyi Biotec GmbH. All rights reserved.