

References for Products 100-118

1. Asquith B, Debacq C, Florins A, Gillet N, Sanchez-Alcaraz T, Mosley A, Willems L. (2006) Quantifying lymphocyte kinetics in vivo using carboxyfluorescein diacetate succinimidyl ester (cfse). *Proc Biol Sci*, 273, 1165.
2. Hashizaki K, Taguchi H, Sakai H, Abe M, Saito Y, Ogawa N. (2006) Carboxyfluorescein leakage from poly(ethylene glycol)-grafted liposomes induced by the interaction with serum. *Chem Pharm Bull (Tokyo)*, 54, 80.
3. Schreer A, Tinson C, Sherry JP, Schirmer K. (2005) Application of alamar blue/5-carboxyfluorescein diacetate acetoxymethyl ester as a noninvasive cell viability assay in primary hepatocytes from rainbow trout. *Anal Biochem*, 344, 76.
4. Wang XQ, Duan XM, Liu LH, Fang YQ, Tan Y. (2005) Carboxyfluorescein diacetate succinimidyl ester fluorescent dye for cell labeling. *Acta Biochim Biophys Sin (Shanghai)*, 37, 379.
5. Tanaka Y, Ohdan H, Onoe T, Mitsuta H, Tashiro H, Itamoto T, Asahara T. (2005) Low incidence of acute rejection after living-donor liver transplantation: Immunologic analyses by mixed lymphocyte reaction using a carboxyfluorescein diacetate succinimidyl ester labeling technique. *Transplantation*, 79, 1262.
6. Pajewski R, Ferdani R, Pajewska J, Djedovic N, Schlesinger PH, Gokel GW. (2005) Evidence for dimer formation by an amphiphilic heptapeptide that mediates chloride and carboxyfluorescein release from liposomes. *Org Biomol Chem*, 3, 619.
7. Watson D, Zhang GY, Sartor M, Alexander SI. (2004) "pruning" of alloreactive cd4+ t cells using 5- (and 6-)carboxyfluorescein diacetate succinimidyl ester prolongs skin allograft survival. *J Immunol*, 173, 6574.
8. Murthy SN, Zhao YL, Sen A, Hui SW. (2004) Cyclodextrin enhanced transdermal delivery of piroxicam and carboxyfluorescein by electroporation. *J Control Release*, 99, 393.
9. Morono Y, Takano S, Miyanaga K, Tanji Y, Unno H, Hori K. (2004) Application of glutaraldehyde for the staining of esterase-active cells with carboxyfluorescein diacetate. *Biotechnol Lett*, 26, 379.
10. Bachmeier CJ, Trickler WJ, Miller DW. (2004) Drug efflux transport properties of 2',7'-bis(2-carboxyethyl)-5(6)-carboxyfluorescein acetoxymethyl ester (bcecf-am) and its fluorescent free acid, bcecf. *J Pharm Sci*, 93, 932.
11. David R, Machova Z, Beck-Sickinger AG. (2003) Semisynthesis and application of carboxyfluorescein-labelled biologically active human interleukin-8. *Biol Chem*, 384, 1619.
12. Seki T, Morimoto K. (2003) Enhancing effects of medium chain aliphatic alcohols and esters on the permeation of 6-carboxyfluorescein and indomethacin through rat skin. *Drug Deliv*, 10, 289.
13. Chen JC, Chang ML, Muench MO. (2003) A kinetic study of the murine mixed lymphocyte reaction by 5,6-carboxyfluorescein diacetate succinimidyl ester labeling. *J Immunol Methods*, 279, 123.
14. Fischer R, Mader O, Jung G, Brock R. (2003) Extending the applicability of carboxyfluorescein in solid-phase synthesis. *Bioconjug Chem*, 14, 653.
15. Mueller-Stahl K, Kofidis T, Akhyari P, Wachsmann B, Lenz A, Boublik J, Heine M, Muehlfait V, Haverich A, Mertsching H. (2003) Carboxyfluorescein diacetate succinimidyl ester facilitates cell tracing and colocalization studies in bioartificial organ engineering. *Int J Artif Organs*, 26, 235.
16. Bergsdorf C, Beyer C, Umansky V, Werr M, Sapp M. (2003) Highly efficient transport of carboxyfluorescein diacetate succinimidyl ester into cos7 cells using human papillomavirus-like particles. *FEBS Lett*, 536, 120.
17. Hoefel D, Grooby WL, Monis PT, Andrews S, Saint CP. (2003) A comparative study of carboxyfluorescein diacetate and carboxyfluorescein diacetate succinimidyl ester as indicators of bacterial activity. *J Microbiol Methods*, 52, 379.

18. Liao RS, Rennie RP, Talbot JA. (2002) Comparative evaluation of a new fluorescent carboxyfluorescein diacetate-modified microdilution method for antifungal susceptibility testing of candida albicans isolates. *Antimicrob Agents Chemother*, 46, 3236.
19. Dumitriu IE, Mohr W, Kolowos W, Kern P, Kalden JR, Herrmann M. (2001) 5,6-carboxyfluorescein diacetate succinimidyl ester-labeled apoptotic and necrotic as well as detergent-treated cells can be traced in composite cell samples. *Anal Biochem*, 299, 247.
20. Yefimov S, Yergey AL, Chrambach A. (2001) Sequential electroelution and mass spectroscopic identification of intact sodium dodecyl sulfate-proteins labeled with 5(6)-carboxyfluorescein-n-hydroxysuccinimide ester. *Electrophoresis*, 22, 2881.
21. Engelk M, Bojarski P, Bloss R, Diehl H. (2001) Tamoxifen perturbs lipid bilayer order and permeability: Comparison of dsc, fluorescence anisotropy, laurdan generalized polarization and carboxyfluorescein leakage studies. *Biophys Chem*, 90, 157.
22. Sathiyaseelan T, Baldwin CL. (2000) Evaluation of cell replication by bovine t cells in polyclonally activated cultures using carboxyfluorescein succinimidyl ester (cfse) loading and flow cytometric analysis. *Res Vet Sci*, 69, 275.
23. Lyons AB, Hasbold J, Hodgkin PD. (2001) Flow cytometric analysis of cell division history using dilution of carboxyfluorescein diacetate succinimidyl ester, a stably integrated fluorescent probe. *Methods Cell Biol*, 63, 375.
24. Lanz E, Slavik J, Kotyk A. (1999) 2',7'-bis-(2-carboxyethyl)-5(6)-carboxyfluorescein as a dual-emission fluorescent indicator of intracellular ph suitable for argon laser confocal microscopy. *Folia Microbiol (Praha)*, 44, 429.
25. Gruber HE, Leslie KP, Ingram JA, Hanley EN, Jr. (2000) Optimization of 5-(and-6)-carboxyfluorescein diacetate succinimidyl ester for labeling human intervertebral disc cells in vitro. *Biotech Histochem*, 75, 118.
26. Massou S, Albigot R, Prats M. (2000) Carboxyfluorescein fluorescence experiments. *Biochem. Educ.*, 28, 171.
27. Hester JD, Lindquist HD, Bobst AM, Schaefer FW, 3rd. (2000) Fluorescent in situ detection of encephalitozoon hellem spores with a 6-carboxyfluorescein-labeled ribosomal rna-targeted oligonucleotide probe. *J Eukaryot Microbiol*, 47, 299.
28. von Horsten S, Helfritz A, Kuhlmann S, Nave H, Tschernig T, Pabst R, Ben-Eliyahu S, Meyer D, Schmidt RE, Schmitz C. (2000) Stereological quantification of carboxyfluorescein-labeled rat lung metastasis: A new method for the assessment of natural killer cell activity and tumor adhesion in vivo and in situ. *J Immunol Methods*, 239, 25.
29. Sun D, Griffith CM, Hay ED. (2000) Carboxyfluorescein as a marker at both light and electron microscope levels to follow cell lineage in the embryo. *Methods Mol Biol*, 135, 357.
30. Nishijo J, Shiota S, Mazima K, Inoue Y, Mizuno H, Yoshida J. (2000) Interactions of cyclodextrins with dipalmitoyl, distearoyl, and dimyristoyl phosphatidyl choline liposomes. A study by leakage of carboxyfluorescein in inner aqueous phase of unilamellar liposomes. *Chem Pharm Bull (Tokyo)*, 48, 48.
31. Fulcher D, Wong S. (1999) Carboxyfluorescein succinimidyl ester-based proliferative assays for assessment of t cell function in the diagnostic laboratory. *Immunol Cell Biol*, 77, 559.
32. Warren HS. (1999) Using carboxyfluorescein diacetate succinimidyl ester to monitor human nk cell division: Analysis of the effect of activating and inhibitory class i mhc receptors. *Immunol Cell Biol*, 77, 544.
33. Mintern J, Li M, Davey GM, Blanas E, Kurts C, Carbone FR, Heath WR. (1999) The use of carboxyfluorescein diacetate succinimidyl ester to determine the site, duration and cell type responsible for antigen presentation in vivo. *Immunol Cell Biol*, 77, 539.
34. Fazekas de St Groth B, Smith AL, Koh WP, Girgis L, Cook MC, Bertolino P. (1999) Carboxyfluorescein diacetate succinimidyl ester and the virgin lymphocyte: A marriage made in heaven. *Immunol Cell Biol*, 77, 530.
35. Hasbold J, Gett AV, Rush JS, Deenick E, Avery D, Jun J, Hodgkin PD. (1999) Quantitative analysis of lymphocyte differentiation and proliferation in vitro using carboxyfluorescein diacetate succinimidyl ester. *Immunol Cell Biol*, 77, 516.

36. Lyons AB. (1999) Divided we stand: Tracking cell proliferation with carboxyfluorescein diacetate succinimidyl ester. *Immunol Cell Biol*, 77, 509.
37. Marechal X, Mordon S, Devoisselle JM, Begu S, Guery B, Neviere R, Buys B, Dhelin G, Lesage JC, Mathieu D, Chopin C. (1999) In vivo application of intestinal pH measurement using 2',7'-bis(carboxyethyl)-5,6-carboxyfluorescein (bcecf) fluorescence imaging. *Photochem Photobiol*, 70, 813.
38. Pohl R, Thrall RS, Rogers RA, Kramer PA. (1999) Confocal imaging of peripheral regions of intact rat lungs following intratracheal administration of 6-carboxyfluorescein, fitc-insulin, and fitc-dextran. *Pharm Res*, 16, 327.
39. Cardullo RA, Dandala S. (1999) Correcting for artifacts in complex aqueous solutions when using the pH-sensitive dye 2',7'-bis-(2-carboxyethyl)-5-(and -6)carboxyfluorescein. *Anal Biochem*, 267, 351.
40. Angulo R, Fulcher DA. (1998) Measurement of candida-specific blastogenesis: Comparison of carboxyfluorescein succinimidyl ester labelling of T cells, thymidine incorporation, and CD69 expression. *Cytometry*, 34, 143.
41. Scott DA, Docampo R, Benchimol M. (1998) Analysis of the uptake of the fluorescent marker 2',7'-bis-(2-carboxyethyl)-5-(and-6)-carboxyfluorescein (bcecf) by hydrogenosomes in *Trichomonas vaginalis*. *Eur J Cell Biol*, 76, 139.
42. Abe F. (1998) Hydrostatic pressure enhances vital staining with carboxyfluorescein or carboxydichlorofluorescein in *Saccharomyces cerevisiae*: Efficient detection of labeled yeasts by flow cytometry. *Appl Environ Microbiol*, 64, 1139.
43. Cohen-Kashi M, Deutsch M, Tirosh R, Rachmani H, Weinreb A. (1997) Carboxyfluorescein as a fluorescent probe for cytoplasmic effects of lymphocyte stimulation. *Spectrochim Acta A Mol Biomol Spectrosc*, 53A, 1655.
44. Bajoria R, Contractor SF. (1997) Effect of surface charge of small unilamellar liposomes on uptake and transfer of carboxyfluorescein across the perfused human term placenta. *Pediatr Res*, 42, 520.
45. Rossi FM, Kao JP. (1997) Practical method for the multigram separation of the 5- and 6-isomers of carboxyfluorescein. *Bioconjug Chem*, 8, 495.
46. Bajoria R, Contractor SF. (1997) Effect of the size of liposomes on the transfer and uptake of carboxyfluorescein by the perfused human term placenta. *J Pharm Pharmacol*, 49, 675.
47. Adamczyk M, Fishpaugh JR, Heuser KJ. (1997) Preparation of succinimidyl and pentafluorophenyl active esters of 5- and 6-carboxyfluorescein. *Bioconjug Chem*, 8, 253.
48. Kosegarten H, Grolig F, Wieneke J, Wilson G, Hoffmann B. (1997) Differential ammonia-elicited changes of cytosolic pH in root hair cells of rice and maize as monitored by 2',7'-bis-(2-carboxyethyl)-5-(and -6)-carboxyfluorescein-fluorescence ratio. *Plant Physiol*, 113, 451.
49. Hirohashi T, Terasaki T, Shigetoshi M, Sugiyama Y. (1997) In vivo and in vitro evidence for nonrestricted transport of 2',7'-bis(2-carboxyethyl)-5(6)-carboxyfluorescein tetraacetoxymethyl ester at the blood-brain barrier. *J Pharmacol Exp Ther*, 280, 813.
50. Draper MP, Martell RL, Levy SB. (1997) Active efflux of the free acid form of the fluorescent dye 2',7'-bis(2-carboxyethyl)-5(6)-carboxyfluorescein in multidrug-resistance-protein-overexpressing murine and human leukemia cells. *Eur J Biochem*, 243, 219.
51. Fujisaki J, Tokunaga Y, Takahashi T, Murata S, Shimojo F, Hata T. (1996) Physicochemical characterization of bisphosphonic carboxyfluorescein for osteotropic drug delivery. *J Pharm Pharmacol*, 48, 798.
52. Kimura M, Araie M, Koyano S. (1996) Movement of carboxyfluorescein across retinal pigment epithelium-choroid. *Exp Eye Res*, 63, 51.
53. Nett W, Deitmer JW. (1996) Simultaneous measurements of intracellular pH in the leech giant glial cell using 2',7'-bis-(2-carboxyethyl)-5,6-carboxyfluorescein and ion-sensitive microelectrodes. *Biophys J*, 71, 394.
54. Garrido T, Riese HH, Quesada AR, Barbacid MM, Aracil M. (1996) Quantitative assay for cell invasion using the fluorogenic substrate 2',7'-bis(2-carboxyethyl)-5-(and -6)-carboxyfluorescein acetoxymethylester. *Anal Biochem*, 235, 234.

55. Ferdous AJ, Ishida T, Shinohara M, Harashima H, Kiwada H. (1996) Size-dependent release of carboxyfluorescein from cetylmannoside-modified liposomes in human plasma. *Biopharm Drug Dispos*, 17, 145.
56. Jackson VN, Halestrap AP. (1996) The kinetics, substrate, and inhibitor specificity of the monocarboxylate (lactate) transporter of rat liver cells determined using the fluorescent intracellular pH indicator, 2',7'-bis(carboxyethyl)-5(6)-carboxyfluorescein. *J Biol Chem*, 271, 861.
57. Breeuwer P, Drocourt J, Rombouts FM, Abee T. (1996) A novel method for continuous determination of the intracellular pH in bacteria with the internally conjugated fluorescent probe 5 (and 6)-carboxyfluorescein succinimidyl ester. *Appl Environ Microbiol*, 62, 178.
58. Fujisaki J, Tokunaga Y, Sawamoto T, Takahashi T, Kimura S, Shimojo F, Hata T. (1996) Osteotropic drug delivery system (odds) based on bisphosphonic prodrug. Iii: Pharmacokinetics and targeting characteristics of osteotropic carboxyfluorescein. *J Drug Target*, 4, 117.
59. Davenpeck KL, Chrest FJ, Sterbinsky SA, Bickel CA, Bochner BS. (1995) Carboxyfluorescein diacetate labeling does not affect adhesion molecule expression or function in human neutrophils or eosinophils. *J Immunol Methods*, 188, 79.
60. Ciray HN, Persson BE, Roomans GM, Ulmsten U. (1995) Dye-coupling between term pregnant human myometrial cells before labor: Carboxyfluorescein versus lucifer yellow. *Cell Biol Int*, 19, 609.
61. Russell DA, Pottier RH, Valenzano DP. (1995) In vivo spectroscopic properties of the fluorescent pH indicator biscarboxyethyl carboxyfluorescein. *J Photochem Photobiol B*, 29, 17.
62. Best L, Elliott AC. (1995) Changes in 2',7'-bis(carboxyethyl) 5'(6')-carboxyfluorescein-, Fura-2 and autofluorescence in intact rat pancreatic islets in response to nutrients and non-nutrients. *Mol Cell Endocrinol*, 111, 191.
63. Breeuwer P, Drocourt JL, Bunschoten N, Zwietering MH, Rombouts FM, Abee T. (1995) Characterization of uptake and hydrolysis of fluorescein diacetate and carboxyfluorescein diacetate by intracellular esterases in *saccharomyces cerevisiae*, which result in accumulation of fluorescent product. *Appl Environ Microbiol*, 61, 1614.
64. Fujisaki J, Tokunaga Y, Takahashi T, Hirose T, Shimojo F, Kagayama A, Hata T. (1995) Osteotropic drug delivery system (odds) based on bisphosphonic prodrug. I: Synthesis and in vivo characterization of osteotropic carboxyfluorescein. *J Drug Target*, 3, 273.
65. Walker B, Gray J, Burns DM, Wang Q, Adrian TE, Nichols DH, Murphy RF, Nelson J. (1995) Carboxyfluorescein and biotin neuromedin c analogues: Synthesis and applications. *Peptides*, 16, 255.
66. Schreiber R, Stoll B, Lang F, Haussinger D. (1994) Effects of aniso-osmolarity and hydroperoxides on intracellular pH in isolated rat hepatocytes as assessed by (2',7')-bis(carboxyethyl)-5(6)-carboxyfluorescein and fluorescein isothiocyanate-dextran fluorescence. *Biochem J*, 303 (Pt 1), 113.
67. Mordon S, Devoisselle JM, Maunoury V. (1994) In vivo pH measurement and imaging of tumor tissue using a pH-sensitive fluorescent probe (5,6-carboxyfluorescein): Instrumental and experimental studies. *Photochem Photobiol*, 60, 274.
68. Fujioka H, Hunt PJ, Rozga J, Wu GD, Cramer DV, Demetriou AA, Moscioni AD. (1994) Carboxyfluorescein (cfse) labelling of hepatocytes for short-term localization following intraportal transplantation. *Cell Transplant*, 3, 397.
69. Hanthamrongwit M, Reid WH, Courtney JM, Grant MH. (1994) 5-carboxyfluorescein diacetate as a probe for measuring the growth of keratinocytes. *Hum Exp Toxicol*, 13, 423.
70. Breeuwer P, Drocourt JL, Rombouts FM, Abee T. (1994) Energy-dependent, carrier-mediated extrusion of carboxyfluorescein from *saccharomyces cerevisiae* allows rapid assessment of cell viability by flow cytometry. *Appl Environ Microbiol*, 60, 1467.
71. Volet B, Lacotte D, Moosmayer M, Anner BM. (1994) Na,k-atpase and carboxyfluorescein distinctly alter vesicle formation in vitro. *Biochim Biophys Acta*, 1191, 1.

72. Petroski RE, Geller HM. (1994) Selective labeling of embryonic neurons cultured on astrocyte monolayers with 5(6)-carboxyfluorescein diacetate (cfda). *J Neurosci Methods*, 52, 23.
73. Kondo M, Araie M. (1994) Movement of carboxyfluorescein across the isolated rabbit iris-ciliary body. *Curr Eye Res*, 13, 251.
74. Ubels JL, Williams KK, Lopez Bernal D, Edelhauser HF. (1994) Evaluation of effects of a physiologic artificial tear on the corneal epithelial barrier: Electrical resistance and carboxyfluorescein permeability. *Adv Exp Med Biol*, 350, 441.
75. Wiedmann TS, Salmon A, Perkins WR. (1993) Comparison of the leakage of carboxyfluorescein from symmetric- and asymmetric-acyl chain phosphatidylcholine vesicles. *Pharm Res*, 10, 147.
76. Griffith CM, Hay ED. (1992) Epithelial-mesenchymal transformation during palatal fusion: Carboxyfluorescein traces cells at light and electron microscopic levels. *Development*, 116, 1087.
77. Provinciali M, Di Stefano G, Fabris N. (1992) Optimization of cytotoxic assay by target cell retention of the fluorescent dye carboxyfluorescein diacetate (cfda) and comparison with conventional 51cr release assay. *J Immunol Methods*, 155, 19.
78. Collington GK, Hunter J, Allen CN, Simmons NL, Hirst BH. (1992) Polarized efflux of 2',7'-bis(2-carboxyethyl)-5(6)-carboxyfluorescein from cultured epithelial cell monolayers. *Biochem Pharmacol*, 44, 417.
79. Owen CS. (1992) Comparison of spectrum-shifting intracellular ph probes 5'(and 6')-carboxy-10-dimethylamino-3-hydroxyspiro[7h-benzo[c]xanthene-7, 1'(3'h)-isobenzofuran]-3'-one and 2',7'-biscarboxyethyl-5(and 6)-carboxyfluorescein. *Anal Biochem*, 204, 65.
80. Mordon S, Maunoury V, Devoisselle JM, Abbas Y, Coustaud D. (1992) Characterization of tumorous and normal tissue using a ph-sensitive fluorescence indicator (5,6-carboxyfluorescein) in vivo. *J Photochem Photobiol B*, 13, 307.
81. Abuknesha RA, al-Mazeedi HM, Price RG. (1992) Reduction of the rate of fluorescence decay of fitc- and carboxyfluorescein-stained cells by anti-fitc antibodies. *Histochem J*, 24, 73.
82. Nagamachi E, Hirai Y, Tomochika K, Kanemasa Y. (1992) Studies on osmotic stability of liposomes prepared with bacterial membrane lipids by carboxyfluorescein release. *Microbiol Immunol*, 36, 231.
83. Tatsu Y, Yamamura S, Yoshikawa S. (1992) Fluorescent fibre-optic immunosensing system based on complement lysis of liposome containing carboxyfluorescein. *Biosens Bioelectron*, 7, 741.
84. Kemenes G, Daykin K, Elliott CJ. (1992) Vital staining with 5(6)-carboxyfluorescein in the feeding system of lymnaea and the selective photoinactivation of the filled neurones. *Acta Biol Hung*, 43, 351.
85. Collington GK, Allen CN, Simmons NL, Hirst BH. (1991) Pharmacological profile of inhibition of 2',7'-bis(2-carboxyethyl)-5(6)-carboxyfluorescein efflux in human hct-8 intestinal epithelial cells. *Biochem Pharmacol*, 42 Suppl, S33.
86. Kemenes G, Daykin K, Elliott CJ. (1991) Photoinactivation of neurones axonally filled with the fluorescent dye 5(6)-carboxyfluorescein in the pond snail, lymnaea stagnalis. *J Neurosci Methods*, 39, 207.
87. Hirnle E, Hirnle P, Wright JK. (1991) Distribution of liposome-incorporated carboxyfluorescein in rabbit eyes. *J Microencapsul*, 8, 391.
88. Saalman E, Norden B, Arvidsson L, Hamnerius Y, Hojevik P, Connell KE, Kurucsev T. (1991) Effect of 2.45 ghz microwave radiation on permeability of unilamellar liposomes to 5(6)-carboxyfluorescein. Evidence of non-thermal leakage. *Biochim Biophys Acta*, 1064, 124.
89. Suzuki Y, Yoshikawa K, Yokochi T. (1991) A new sensitive and rapid automated fluorometric assay for detection of natural killer activity using carboxyfluorescein diacetate. *J Immunoassay*, 12, 145.

90. Allen CN, Harpur ES, Gray TJ, Simmons NL, Hirst BH. (1990) Efflux of bis-carboxyethyl-carboxyfluorescein (bcecf) by a novel atp-dependent transport mechanism in epithelial cells. *Biochem Biophys Res Commun*, 172, 262.
91. Fang T, Naguib KS, Peyman GA, Khoobei B. (1990) Comparative study of three fluorescent dyes for angiography: Sodium fluorescein, carboxyfluorescein, and calcein. *Ophthalmic Surg*, 21, 250.
92. Domingo JC, Rosell F, Mora M, de Madariaga MA. (1989) Importance of the purification grade of 5(6)-carboxyfluorescein on the stability and permeability properties of n-acylphosphatidylethanolamine liposomes. *Biochem Soc Trans*, 17, 997.
93. Miyata H, Hayashi H, Suzuki S, Noda N, Kobayashi A, Fujiwake H, Hirano M, Yamazaki N. (1989) Dual loading of the fluorescent indicator fura-2 and 2,7-bis(carboxyethyl)-5(6)-carboxyfluorescein (bcecf) in isolated myocytes. *Biochem Biophys Res Commun*, 163, 500.
94. Tucker JE, Mauzerall D, Tucker EB. (1989) Symplastic transport of carboxyfluorescein in staminal hairs of *Setcreasea purpurea* is diffusive and includes loss to the vacuole. *Plant Physiol*, 90, 1143.
95. Liu ZY, Solow R, Hu VW. (1988) Fluorescence analysis of size distribution and mode of dye release from carboxyfluorescein-loaded vesicles: Application to the study of complement-membrane interactions. *Biochim Biophys Acta*, 945, 253.
96. Grimes PA. (1988) Carboxyfluorescein distribution in ocular tissues of normal and diabetic rats. *Curr Eye Res*, 7, 981.
97. Tsujimoto K, Semadeni M, Huflejt M, Packer L. (1988) Intracellular ph of halobacteria can be determined by the fluorescent dye 2', 7'-bis(carboxyethyl)-5(6)-carboxyfluorescein. *Biochem Biophys Res Commun*, 155, 123.
98. Chen RF, Knutson JR. (1988) Mechanism of fluorescence concentration quenching of carboxyfluorescein in liposomes: Energy transfer to nonfluorescent dimers. *Anal Biochem*, 172, 61.
99. Grimes PA. (1988) Carboxyfluorescein transfer across the blood-retinal barrier evaluated by quantitative fluorescence microscopy: Comparison with fluorescein. *Exp Eye Res*, 46, 769.
100. Kolber MA, Quinones RR, Gress RE, Henkart PA. (1988) Measurement of cytotoxicity by target cell release and retention of the fluorescent dye bis-carboxyethyl-carboxyfluorescein (bcecf). *J Immunol Methods*, 108, 255.
101. Davis MH, Altschuld RA, Jung DW, Brierley GP. (1987) Estimation of intramitochondrial pca and ph by fura-2 and 2,7 bis(carboxyethyl)-5(6)-carboxyfluorescein (bcecf) fluorescence. *Biochem Biophys Res Commun*, 149, 40.
102. Bramhall J, Hofmann J, DeGuzman R, Montestruque S, Schell R. (1987) Temperature dependence of membrane ion conductance analyzed by using the amphiphilic anion 5/6-carboxyfluorescein. *Biochemistry*, 26, 6330.
103. Touitou E, Alhaique F, Fisher P, Memoli A, Riccieri FM, Santucci E. (1987) Prevention of molecular self-association by sodium salicylate: Effect on insulin and 6-carboxyfluorescein. *J Pharm Sci*, 76, 791.
104. Au S, Weiner ND, Schacht J. (1987) Aminoglycoside antibiotics preferentially increase permeability in phosphoinositide-containing membranes: A study with carboxyfluorescein in liposomes. *Biochim Biophys Acta*, 902, 80.
105. Hansson Y, Jacobson E, Ortlund J, Paulie S, Perlmann P. (1987) A rapid method for detection of cellular proliferation using carboxyfluorescein. Assay of growth factors (il-2, il-1) and growth inhibiting antibodies. *J Immunol Methods*, 100, 261.
106. Tsuboi S, Pederson JE. (1987) Permeability of the blood-retinal barrier to carboxyfluorescein in eyes with rhegmatogenous retinal detachment. *Invest Ophthalmol Vis Sci*, 28, 96.
107. Tsuboi S, Pederson JE. (1987) Acetazolamide effect on the inward permeability of the blood-retinal barrier to carboxyfluorescein. *Invest Ophthalmol Vis Sci*, 28, 92.
108. Tsuboi S, Pederson JE. (1986) Permeability of the isolated dog retinal pigment epithelium to carboxyfluorescein. *Invest Ophthalmol Vis Sci*, 27, 1767.

109. Barber RF, Shek PN. (1986) Liposomes and tear fluid. I. Release of vesicle-entrapped carboxyfluorescein. *Biochim Biophys Acta*, 879, 157.
110. Kayalar C, Duzgunes N. (1986) Membrane action of colicin e1: Detection by the release of carboxyfluorescein and calcein from liposomes. *Biochim Biophys Acta*, 860, 51.
111. Papa S, Caramelli E, Allegri G, Billi AM, Manzoli FA, Capitani S. (1986) Flow cytometric analysis of liposome-nuclei interaction: Transfer and intranuclear release of carboxyfluorescein. *Cytometry*, 7, 331.
112. Rao G, Barnes CA, McNaughton BL. (1986) Intracellular fluorescent staining with carboxyfluorescein: A rapid and reliable method for quantifying dye-coupling in mammalian central nervous system. *J Neurosci Methods*, 16, 251.
113. Verkman AS, Ives HE. (1986) Anomalous driving force for renal brush border h⁺/oh⁻-transport characterized by using 6-carboxyfluorescein. *Biochemistry*, 25, 2876.
114. Neault TR, McLaren JW, Brubaker JH, Brubaker RF. (1986) Spectral shift of fluorescein and carboxyfluorescein in the anterior chamber of the rabbit eye following systemic administration. *Curr Eye Res*, 5, 337.
115. Woolfrey SG, Taylor G, Kellaway IW, Smith A. (1986) Pulmonary absorption of carboxyfluorescein in the rat. *J Pharm Pharmacol*, 38, 304.
116. Ohshima T, Yoshikawa H, Takada K, Muranishi S. (1986) Enhancing effect of absorption promoters on percutaneous absorption of a model dye (6-carboxyfluorescein) as a poorly absorbable drug. Iii. Histological study after addition of various absorption promoters in rats. *J Pharmacobiodyn*, 9, 223.
117. Araie M. (1986) Carboxyfluorescein. A dye for evaluating the corneal endothelial barrier function in vivo. *Exp Eye Res*, 42, 141.
118. Weinstein JN, Blumenthal R, Klausner RD. (1986) Carboxyfluorescein leakage assay for lipoprotein-liposome interaction. *Methods Enzymol*, 128, 657.
119. Blair NP, Rusin MM. (1986) Blood-retinal barrier permeability to carboxyfluorescein and fluorescein in monkeys. *Graefes Arch Clin Exp Ophthalmol*, 224, 419.
120. Ohshima T, Yoshikawa H, Takada K, Muranishi S. (1985) Enhancing effect of absorption promoters on percutaneous absorption of a model dye (6-carboxyfluorescein) as poorly absorbable drugs. Ii. Study on the absorption promoting effect of azone. *J Pharmacobiodyn*, 8, 900.
121. Amselem S, Loyter A, Lichtenberg D, Barenholz Y. (1985) The interaction of sendai virus with negatively charged liposomes: Virus-induced lysis of carboxyfluorescein-loaded small unilamellar vesicles. *Biochim Biophys Acta*, 820, 1.
122. Mornet D, Ue K. (1985) Incorporation of 6-carboxyfluorescein into myosin subfragment 1. *Biochemistry*, 24, 840.
123. Blair NP, Jones CW, Rusin MM. (1984) Pathophysiology of the blood-retinal barrier in experimental diabetes. Vitreous fluorophotometry using carboxyfluorescein and fluorescein. *Arch Ophthalmol*, 102, 1810.
124. Stevenson M, Baillie AJ, Richards RM. (1984) Quantification of uptake of liposomal carboxyfluorescein by professional phagocytes in-vitro. A flow microfluorimetric study on the j774 murine macrophage cell line. *J Pharm Pharmacol*, 36, 824.
125. Ohshima T, Yoshikawa H, Takada K, Muranishi S. (1984) Enhancing effect of absorption promoters on percutaneous absorption of a model dye (6-carboxyfluorescein) as poorly absorbable drugs. I. Comparison of plasma levels after addition of various absorption promoters in rat. *J Pharmacobiodyn*, 7, 648.
126. Barbet J, Machy P, Truneh A, Leserman LD. (1984) Weak acid-induced release of liposome-encapsulated carboxyfluorescein. *Biochim Biophys Acta*, 772, 347.
127. Hashida N, Murakami M, Yoshikawa H, Takada K, Muranishi S. (1984) Intestinal absorption of carboxyfluorescein entrapped in liposomes in comparison with its administration with lipid-surfactant mixed micelles. *J Pharmacobiodyn*, 7, 195.
128. Ogura Y, Tsukahara Y, Saito I, Kondo T. (1984) [fluorophotometry with carboxyfluorescein]. *Nippon Ganka Gakkai Zasshi*, 88, 230.
129. Papa S, Caramelli E, Chiarini C, Allegri G, Capitani S. (1984) Transfer of liposome-encapsulated carboxyfluorescein to isolated nuclei. *Cell Biol Int Rep*, 8, 27.

130. Rockey JH, Li W, Eccleston JF. (1983) Binding of fluorescein and carboxyfluorescein by human serum proteins: Significance of kinetic and equilibrium parameters of association in ocular fluorometric studies. *Exp Eye Res*, 37, 455.
131. Truneh A, Machy P, Barbet J, Mishal Z, Lemonnier FA, Leserman LD. (1983) Endocytosis of hla and h-2 molecules on transformed murine cells measured by fluorescence dequenching of liposome-encapsulated carboxyfluorescein. *Embo J*, 2, 2285.
132. Roozmond RC, Urli DC. (1982) Peculiar behaviour of rabbit thymocytes in interaction with liposomes of different compositions shown by fluorescence polarization studies, lipid analysis, and uptake of vesicle-entrapped carboxyfluorescein. *Biochim Biophys Acta*, 689, 499.
133. Lelkes PI, Tandeter HB. (1982) Studies on the methodology of the carboxyfluorescein assay and on the mechanism of liposome stabilization by red blood cells in vitro. *Biochim Biophys Acta*, 716, 410.
134. Grimes PA, Stone RA, Laties AM, Li W. (1982) Carboxyfluorescein. A probe of the blood-ocular barriers with lower membrane permeability than fluorescein. *Arch Ophthalmol*, 100, 635.
135. Goodall H, Johnson MH. (1982) Use of carboxyfluorescein diacetate to study formation of permeable channels between mouse blastomeres. *Nature*, 295, 524.
136. Rockey JH, Li W. (1982) Binding of fluorescein and carboxyfluorescein by normal and glycosylated human serum proteins. *Ophthalmic Res*, 14, 416.
137. Leserman LD, Weinstein JN, Moore JJ, Terry WD. (1980) Specific interaction of myeloma tumor cells with hapten-bearing liposomes containing methotrexate and carboxyfluorescein. *Cancer Res*, 40, 4768.
138. Khanna PL, Ullman EF. (1980) 4',5'-dimethoxy-6-carboxyfluorescein: A novel dipole-dipole coupled fluorescence energy transfer acceptor useful for fluorescence immunoassays. *Anal Biochem*, 108, 156.
139. Bruning JW, Kardol MJ, Arentzen R. (1980) Carboxyfluorescein fluorochromasia assays. I. Non-radioactively labeled cell mediated lympholysis. *J Immunol Methods*, 33, 33.
140. Bruning JW, Kardol MJ, Arentzen R, Naipal A, van der Poel JJ. (1979) Carboxyfluorescein fluorochromasia assays for cell-mediated lympholysis (cml) and antibody-dependent cellular cytotoxicity (adcc): A nonradioactive technique. *Transplant Proc*, 11, 1961.