

## **References for Product 14003**

1. Warnakulasuriyarachchi D, Ungureanu NH, Holcik M. (2003) The translation of an antiapoptotic protein HIAP2 is regulated by an upstream open reading frame. *Cell Death Differ*, 10, 899.
2. Cheng L, Goldman E. (2001) Absence of effect of varying Thr-Leu codon pairs on protein synthesis in a T7 system. *Biochemistry*, 40, 6102.
3. Takahashi N, Sasagawa N, Suzuki K, Ishiura S. (2000) The CUG-binding protein binds specifically to UG dinucleotide repeats in a yeast three-hybrid system. *Biochem Biophys Res Commun*, 277, 518.
4. Van Etten WJ, Janssen GR. (1998) An AUG initiation codon, not codon-anticodon complementarity, is required for the translation of unleadered mRNA in *Escherichia coli*. *Mol Microbiol*, 27, 987.
5. Antoine M, Reimers K, Dickson C, Kiefer P. (1997) Fibroblast growth factor 3, a protein with dual subcellular localization, is targeted to the nucleus and nucleolus by the concerted action of two nuclear localization signals and a nucleolar retention signal. *J Biol Chem*, 272, 29475.
6. Babic S, Hunter CN, Rakhlin NJ, Simons RW, Phillips-Jones MK. (1997) Molecular characterisation of the pifC gene encoding translation initiation factor 3, which is required for normal photosynthetic complex formation in *Rhodobacter sphaeroides* NCIB 8253. *Eur J Biochem*, 249, 564.
7. Shi J, Friedman S, Maciag T. (1997) A carboxyl-terminal domain in fibroblast growth factor (FGF)-2 inhibits FGF-1 release in response to heat shock in vitro. *J Biol Chem*, 272, 1142.
8. Siple J, Goldman E. (1993) Increased ribosomal accuracy increases a programmed translational frameshift in *Escherichia coli*. *Proc Natl Acad Sci U S A*, 90, 2315.
9. Khudyakov Yu E, Neplyueva VS, Kalinina TI, Smirnov VD. (1988) Effect of structure of the initiator codon on translation in *E. coli*. *FEBS Lett*, 232, 369.
10. Miller JH, Albertini AM. (1983) Effects of surrounding sequence on the suppression of nonsense codons. *J Mol Biol*, 164, 59.