

## References for Product 17030

1. Zeng Y, Wang Y. (2006) Sequence-dependent formation of intrastrand crosslink products from the UVB irradiation of duplex DNA containing a 5-bromo-2'-deoxyuridine or 5-bromo-2'-deoxycytidine. *Nucleic Acids Res.*
2. Kuwagata M, Ogawa T, Nagata T, Shioda S. (2006) The evaluation of early embryonic neurogenesis after exposure to the genotoxic agent 5-bromo-2'-deoxyuridine in mice. *Neurotoxicology.*
3. Sahambi SK, Hales BF. (2006) Exposure to 5-bromo-2'-deoxyuridine induces oxidative stress and activator protein-1 DNA binding activity in the embryo. *Birth Defects Res A Clin Mol Teratol*, 76, 580.
4. Hoyk Z, Varga C, Parducz A. (2006) Estrogen-induced region specific decrease in the density of 5-bromo-2'-deoxyuridine-labeled cells in the olfactory bulb of adult female rats. *Neuroscience*, 141, 1919.
5. Wojcik A, Bochenek A, Lankoff A, Lisowska H, Padjas A, Szumiel I, von Sonntag C, Obe G. (2006) DNA interstrand crosslinks are induced in cells prelabelled with 5-bromo-2'-deoxyuridine and exposed to UVC radiation. *J Photochem Photobiol B*, 84, 15.
6. Caldwell MA, He X, Svendsen CN. (2005) 5-Bromo-2'-deoxyuridine is selectively toxic to neuronal precursors in vitro. *Eur J Neurosci*, 22, 2965.
7. Yamada K, Semba R, Ding X, Ma N, Nagahama M. (2005) Discrimination of cell nuclei in early S-phase, mid-to-late S-phase, and G(2)/M-phase by sequential administration of 5-bromo-2'-deoxyuridine and 5-chloro-2'-deoxyuridine. *J Histochem Cytochem*, 53, 1365.
8. Shih CH, Whalen BA, Goto Y, Hogg JC, van Eeden SF. (2005) Flow cytometric method for enumeration and characterization of newly released polymorphonuclear leukocytes from the bone marrow using 5'-bromo-2'-deoxyuridine. *Am J Physiol Cell Physiol*, 289, C757.
9. Ogawa T, Kuwagata M, Muneoka KT, Shioda S. (2005) Neuropathological examination of fetal rat brain in the 5-bromo-2'-deoxyuridine-induced neurodevelopmental disorder model. *Congenit Anom (Kyoto)*, 45, 14.
10. Kuwagata M, Muneoka KT, Ogawa T, Takigawa M, Nagao T. (2004) Locomotor hyperactivity following prenatal exposure to 5-bromo-2'-deoxyuridine: neurochemical and behavioral evidence of dopaminergic and serotonergic alterations. *Toxicol Lett*, 152, 63.
11. Ma XD, Ke T, Li YX, Huang CX, Song AD, Chen HG, Wu YH, Jia XC, He GY. (2004) In vitro mutagenesis of *Xanthomonas campestris* alpha-amylase gene by partially replacing deoxythymidine triphosphate with 5-bromo-2'-deoxyuridine-5'-triphosphate using a PCR technique. *Biotechnol Lett*, 26, 171.
12. Ito T, Rokita SE. (2003) Excess electron transfer from an internally conjugated aromatic amine to 5-bromo-2'-deoxyuridine in DNA. *J Am Chem Soc*, 125, 11480.
13. De Boer RJ, Mohri H, Ho DD, Perelson AS. (2003) Estimating average cellular turnover from 5-bromo-2'-deoxyuridine (BrdU) measurements. *Proc Biol Sci*, 270, 849.
14. Goto Y, Hogg JC, Suwa T, Quinlan KB, van Eeden SF. (2003) A novel method to quantify the turnover and release of monocytes from the bone marrow using the thymidine analog 5'-bromo-2'-deoxyuridine. *Am J Physiol Cell Physiol*, 285, C253.
15. Gauthier DT, Cartwright DD, Densmore CL, Blazer VS, Ottinger CA. (2003) Measurement of in vitro leucocyte mitogenesis in fish: ELISA based detection of the thymidine analogue 5'-bromo-2'-deoxyuridine. *Fish Shellfish Immunol*, 14, 279.
16. Wojcik A, von Sonntag C, Obe G. (2003) Application of the biotin-dUTP chromosome labelling technique to study the role of 5-bromo-2'-deoxyuridine in the formation of UV-induced sister chromatid exchanges in CHO cells. *J Photochem Photobiol B*, 69, 139.
17. Tamarit A, Alberola J, Galindo MJ, Navarro D. (2002) Application of a 5-bromo-2'-deoxyuridine ELISA for measuring the lymphoproliferative response to human cytomegalovirus in HIV-1-infected patients. *J Virol Methods*, 105, 247.

18. Fujii M, Ito H, Hasegawa T, Suzuki T, Adachi N, Ayusawa D. (2002) 5-Bromo-2'-deoxyuridine efficiently suppresses division potential of the yeast *Saccharomyces cerevisiae*. *Biosci Biotechnol Biochem*, 66, 906.
19. Butov AA, Volkov MA, Anisimov VN. (2001) [Mathematical and simulating model of accelerated aging induced by 5-bromo-2'-deoxyuridine]. *Adv Gerontol*, 8, 70.
20. Brennan LM, Boncavage-Hennessey EM, Wolfe MJ, Toussaint MW, Dennis WE, Rosencrance AB, Gardner HS, Jr. (2001) An in vivo method for using 5-bromo-2'-deoxyuridine (BrdU) as a marker of chemically-induced hepatocellular proliferation in the Japanese medaka (*Oryzias latipes*). *Toxicol Pathol*, 29, 387.
21. Takeyoshi M, Yamasaki K, Yakabe Y, Takatsuki M, Kimber I. (2001) Development of non-radio isotopic endpoint of murine local lymph node assay based on 5-bromo-2'-deoxyuridine (BrdU) incorporation. *Toxicol Lett*, 119, 203.
22. Ohta Y, Ichimura K. (2000) Proliferation markers, proliferating cell nuclear antigen, Ki67, 5-bromo-2'-deoxyuridine, and cyclin D1 in mouse olfactory epithelium. *Ann Otol Rhinol Laryngol*, 109, 1046.
23. Zhivkova Z, Stankova I. (2000) Stability of some novel thymidine, 5-bromo-2'-deoxyuridine and 3'-azido-2'-3'-dideoxythymidine analogues. *Int J Pharm*, 200, 181.
24. Bonhoeffer S, Mohri H, Ho D, Perelson AS. (2000) Quantification of cell turnover kinetics using 5-bromo-2'-deoxyuridine. *J Immunol*, 164, 5049.
25. Struikmans H, Rutgers DH, Jansen GH, Dullens HF, Oosten L, Tulleken CA, van der Tweel I, Battermann JJ. (1999) Prognostic relevance of MIB-1 immunoreactivity, S-phase fraction, 5-bromo-2'-deoxyuridine labeling indices, and mitotic figures in gliomas. *Radiat Oncol Investig*, 7, 243.
26. Biegel LB, Applegate M, Sulecki LM. (1999) Effects of 5-bromo-2'-deoxyuridine (BrdU) implants on hepatic cytochrome P-450 content and beta-oxidation activity in rats and mice. *Drug Chem Toxicol*, 22, 481.
27. Mancini WR, Glaze ER, Stetson PL, Greenberg HS. (1999) Sensitization of 1,3-bis(2-chloroethyl)-1-nitrosourea and cisplatin cytotoxicity by 5-bromo-2'-deoxyuridine in human glioma. *J Pharmacol Exp Ther*, 289, 1404.
28. Maghni K, Nicolescu OM, Martin JG. (1999) Suitability of cell metabolic colorimetric assays for assessment of CD4+ T cell proliferation: comparison to 5-bromo-2'-deoxyuridine (BrdU) ELISA. *J Immunol Methods*, 223, 185.
29. Kuwagata M, Nagao T. (1998) Behavior and reproductive function of rat male offspring treated prenatally with 5-bromo-2'-deoxyuridine. *Reprod Toxicol*, 12, 541.
30. Nagao T, Kuwagata M, Saito Y. (1998) Effects of prenatal exposure to 5-bromo-2'-deoxyuridine on the developing brain and reproductive function in male mouse offspring. *Reprod Toxicol*, 12, 477.
31. Potapenko AI, Rudakovskaia EG, Kozhevnikova NA, Putrina ID, Akif'ev AP. (1997) [The effect of 5-bromo-2'-deoxyuridine on the lifespan and behavior of *Drosophila melanogaster*]. *Ontogenez*, 28, 463.
32. Kiliass D, Macrae FA, Sharpe K, Young GP. (1997) The effect of incubation of rectal biopsies on measures of proliferation using proliferating cell nuclear antigen in comparison with 5-bromo-2'-deoxyuridine. *Cancer Epidemiol Biomarkers Prev*, 6, 819.
33. Nagao T, Sato M, Kuwagata M, Saito Y. (1997) Effects of prenatal exposure to 5-bromo-2'-deoxyuridine on reproductive function in male mouse offspring. *Reprod Toxicol*, 11, 663.
34. Steinberg JJ, Oliver GW, Jr., Farah N, Simoni P, Winiarsky R, Cajigas A. (1997) In vivo determination of 5-bromo-2'-deoxyuridine incorporation into DNA tumor tissue by a new <sup>32</sup>P-postlabelling thin-layer chromatographic method. *J Chromatogr B Biomed Sci Appl*, 694, 333.
35. Akif'ev AP, Potapenko AI, Rudakovskaia EG. (1997) [Ionizing radiation and 5-bromo-2'-deoxyuridine as tools for analysis of the fundamental mechanism of aging in animals]. *Radiats Biol Radioecol*, 37, 613.
36. Kawabe M, Shibata MA, Sano M, Takesada Y, Tamano S, Ito N, Shirai T. (1997) Decrease of prostaglandin E2 and 5-bromo-2'-deoxyuridine labeling but not prostate

- tumor development by indomethacin treatment of rats given 3,2'-dimethyl-4-aminobiphenyl and testosterone propionate. *Jpn J Cancer Res*, 88, 350.
37. Jecker P, Beuleke A, Dressendorfer I, Pabst R, Westermann J. (1997) Long-term oral application of 5-bromo-2'-deoxyuridine does not reliably label proliferating immune cells in the LEW rat. *J Histochem Cytochem*, 45, 393.
  38. de Roos WK, von Geusau BA, Bouwman E, van Dierendonck JH, Borel Rinkes IH, Terpstra OT. (1997) Monitoring engraftment of transplanted hepatocytes in recipient liver with 5-bromo-2'-deoxyuridine. *Transplantation*, 63, 513.
  39. Malendowicz LK, Rebuffat P, Andreis PG, Nussdorfer GG, Nowak M. (1997) Different mechanisms mediate the in vivo aldosterone and corticosterone responses to 5-bromo-2'-deoxyuridine in rats. *Exp Clin Endocrinol Diabetes*, 105, 277.
  40. Menne S, Maschke J, Tolle T, Kreuzfelder E, Grosse-Wilde H, Roggendorf M. (1997) Determination of peripheral blood mononuclear cell responses to mitogens and woodchuck hepatitis virus core antigen in woodchucks by 5-bromo-2'-deoxyuridine or 2[3H]adenine incorporation. *Arch Virol*, 142, 511.
  41. Malendowicz LK, Nussdorfer GG, Trejer M. (1997) Effect of 5-bromo-2'-deoxyuridine on the proliferative activity of thymus and regenerating adrenal cortex. *Life Sci*, 61, 641.
  42. Uporov AV, Tsyrlina EV, Semiglazov VF, Pozharisskii KM. (1997) [Determination of the proliferative activity of breast cancer cells using 5-bromo-2'-deoxyuridine in vivo]. *Vopr Onkol*, 43, 176.
  43. Nagao T, Saito Y, Watanabe C, Imai K. (1997) Reproductive effects of prenatal exposure to 5-bromo-2'-deoxyuridine on male rats. *Reprod Toxicol*, 11, 37.
  44. Miyagawa M, Katsuta O, Tsuchitani M, Yoshikawa K. (1997) Measurement of replicative DNA synthesis (RDS) by a 5-bromo-2'-deoxyuridine (BrdU) labeling technique for detection of hepatocyte proliferation. *J Vet Med Sci*, 59, 45.
  45. Bourgeois CA, Dennebouy R, Gibaud A, Gerbault-Seureau M, Malfroy B, Slodzian G, Galle P, Dutrillaux B. (1996) Scanning ion analytical microscopy for high-resolution detection of 5-bromo-2'-deoxyuridine incorporation in metaphase chromosomes. *Chromosome Res*, 4, 574.
  46. Kuan HY, Smith DE, Ensminger WD, Knol JA, DeRemer SJ, Yang Z, Stetson PL. (1996) Regional pharmacokinetics of 5-bromo-2'-deoxyuridine and 5-fluorouracil in dogs: hepatic arterial versus portal venous infusions. *Cancer Res*, 56, 4724.
  47. Malendowicz LK, Nussdorfer GG. (1996) 5-Bromo-2'-deoxyuridine stimulates the pituitary-adrenal axis in the rat: an effect blocked partially by endothelin-receptor antagonists. *J Int Med Res*, 24, 363.
  48. Muhlmann-Diaz MC, Dullea RG, Bedford JS. (1996) Application of 5-bromo-2'-deoxyuridine as a label for in situ hybridization in chromosome microdissection and painting, and 3' OH DNA end labeling for apoptosis. *Biotechniques*, 21, 82.
  49. Kinsella TJ. (1996) Correspondence re: J.A. Knol et al., Incorporation of 5-bromo-2'-deoxyuridine into colorectal liver metastases and liver in patients receiving a 7-day hepatic arterial infusion. *Cancer Res.*, 55: 3687-3691, 1995. *Cancer Res*, 56, 2677.
  50. Anisimov VN, Sheiko EV. (1996) [Skin carcinogenesis induced by neonatal administration of 5-bromo-2'-deoxyuridine and subsequent treatment with 12-O-tetradecanoylphorbol-13-acetate in mice]. *Vopr Onkol*, 42, 82.
  51. Sawada S, Asakura S, Daimon H, Furihata C. (1995) Comparison of autoradiography, liquid scintillation counting and immunoenzymatic staining of 5-bromo-2'-deoxyuridine for measurement of unscheduled DNA synthesis and replicative DNA synthesis in rat liver. *Mutat Res*, 344, 109.
  52. Knol JA, Walker SC, Robertson JM, Yang Z, DeRemer S, Stetson PL, Ensminger WD, Lawrence TS. (1995) Incorporation of 5-bromo-2'-deoxyuridine into colorectal liver metastases and liver in patients receiving a 7-day hepatic arterial infusion. *Cancer Res*, 55, 3687.
  53. Kitchin KT, Brown JL. (1995) Incorporation of 5-iodo-2'-deoxyuridine and 5-bromo-2'-deoxyuridine into rodent DNA as determined by neutron activation analysis. *Anal Biochem*, 229, 180.

54. Anisimov VN. (1995) Carcinogenesis induced by neonatal exposure to various doses of 5-bromo-2'-deoxyuridine in rats. *Cancer Lett*, 91, 63.
55. Kodama E, Mori S, Shigeta S. (1995) Analysis of mutations in the thymidine kinase gene of varicella zoster virus associated with resistance to 5-iodo-2'-deoxyuridine and 5-bromo-2'-deoxyuridine. *Antiviral Res*, 27, 165.
56. Lerner-Tung MB, Doong SL, Cheng YC, Hsiung GD. (1995) Characterization of conditions for the activation of endogenous guinea pig retrovirus in cultured cells by 5-bromo-2'-deoxyuridine. *Virus Genes*, 9, 201.