

References for Products 21050 and 21052

1. Goko H, Matsuoka A. (1999) W-5 and quin 2-AM reverse the inhibitory effect of insulin on lipolysis due to dibutyryl cAMP. *Diabetes Res Clin Pract*, 44, 101.
2. Faux SP, Michelangeli F, Levy LS. (1994) Calcium chelator Quin-2 prevents crocidolite-induced DNA strand breakage in human white blood cells. *Mutat Res*, 311, 209.
3. Lakowicz JR, Szmajcinski H, Nowaczyk K, Lederer WJ, Kirby MS, Johnson ML. (1994) Fluorescence lifetime imaging of intracellular calcium in COS cells using Quin-2. *Cell Calcium*, 15, 7.
4. Rao GH, Gerrard JM, Murthy M, White JG. (1993) Possible mechanisms of epinephrine actions in quin-2-loaded platelets refractory to arachidonic acid. *Biochem Med Metab Biol*, 50, 322.
5. Lakowicz JR, Szmajcinski H, Nowaczyk K, Johnson ML. (1992) Fluorescence lifetime imaging of calcium using Quin-2. *Cell Calcium*, 13, 131.
6. Carpenter-Deyo L, Reed DJ. (1991) Involvement of calcium and iron in Quin 2 toxicity to isolated hepatocytes. *J Pharmacol Exp Ther*, 258, 747.
7. Carpenter-Deyo L, Duimstra JR, Hedstrom O, Reed DJ. (1991) Toxicity to isolated hepatocytes caused by the intracellular calcium indicator, Quin 2. *J Pharmacol Exp Ther*, 258, 739.
8. Ruchko MV, Slatvinskaia EA, Matyshevskaja OP, Kucherenko NE. (1990) [Increase in free Ca²⁺ level determined by quin-2 in spleen lymphocytes from irradiated rats]. *Ukr Biokhim Zh*, 62, 110.
9. Jefferson JR, Hunt JB, Ginsburg A. (1990) Characterization of indo-1 and quin-2 as spectroscopic probes for Zn²⁺-protein interactions. *Anal Biochem*, 187, 328.
10. Mazorow DL, Millar DB. (1990) Quin-2 and fura-2 measure calcium differently. *Anal Biochem*, 186, 28.
11. Rao GH, White JG. (1990) Aspirin, prostaglandin E1 and Quin-2 AM-induced platelet dysfunction: restoration of function by noradrenalin. *Prostaglandins Leukot Essent Fatty Acids*, 39, 141.
12. Blanco P, Martinez-Serrano A, Bogonez E, Satrustegui J. (1990) Effect of Quin-2 on Ca²⁺ uptake mediated by Na⁺/Ca²⁺ exchange and ⁴⁵Ca²⁺ efflux in rat brain synaptosomes: a requirement for [Ca²⁺]_i. *Cell Calcium*, 11, 25.
13. Cantoni O, Sestili P, Cattabeni F, Bellomo G, Pou S, Cohen M, Cerutti P. (1989) Calcium chelator Quin 2 prevents hydrogen-peroxide-induced DNA breakage and cytotoxicity. *Eur J Biochem*, 182, 209.
14. Gilroy S, Hughes WA, Trewavas AJ. (1989) A Comparison between Quin-2 and Aequorin as Indicators of Cytoplasmic Calcium Levels in Higher Plant Cell Protoplasts. *Plant Physiol*, 90, 482.
15. Vijayaraghavan S, Hoskins D. (1989) Quantitation of bovine sperm cytoplasmic calcium with Quin-2 and Fura-2: evidence that external calcium does not have direct access to the sperm cytoplasm. *Cell Calcium*, 10, 241.
16. Raspe E, Ramboer I, Galand N, Boeynaems JM. (1989) Enhanced release of prostacyclin from quin 2-loaded endothelial cells. *Eur J Pharmacol*, 163, 345.
17. Komada H, Nakabayashi H, Nakano H, Hara M, Yoshida T, Takanari H, Izutsu K. (1989) Measurement of the cytosolic free calcium ion concentration of individual lymphocytes by microfluorometry using quin 2 or fura-2. *Cell Struct Funct*, 14, 141.
18. Noronha-Blob L, Richard C, U'Prichard DC. (1988) Voltage-sensitive calcium channels in differentiated neuroblastoma X glioma hybrid (NG108-15) cells: characterization by quin 2 fluorescence. *J Neurochem*, 50, 1381.
19. Simpson AM, White IG. (1988) Measurement and manipulation of cytoplasmic free calcium of ram and boar spermatozoa using quin 2. *Cell Calcium*, 9, 45.
20. Suko J, Wyskovsky W, Pidlich J, Plank B, Hauptner R, Hellmann G. (1988) Calcium dissociation from mastoparan complexes of calmodulin and its tryptic fragments measured with Quin 2. *Prog Clin Biol Res*, 252, 161.

21. Moore DC, Stanisstreet M. (1988) Perturbation of in vitro development of rodent embryos by calcium antagonist Quin-2. *Cytobios*, 55, 133.
22. Korc M, Schoni MH. (1988) Quin 2 and manganese define multiple alterations in cellular calcium homeostasis in diabetic rat pancreas. *Diabetes*, 37, 13.
23. de Chaffoy de Courcelles D, Roevens P, Verheyen F, Van Belle H, De Clerck F. (1987) The effect of the intracellular calcium chelator Quin-2 on the platelet phosphoinositide metabolism, protein phosphorylation and morphology. *Thromb Haemost*, 58, 927.
24. Noronha-Blob L, Richard C, U'Prichard DC. (1987) Calcium mobilization by muscarinic receptors in human astrocytoma cells: measurements with quin 2. *Biochem Biophys Res Commun*, 147, 182.
25. Maruyama I, Oyamada H, Hasegawa T, Ohtsuka K, Momose K. (1987) Effect of DFP on loading of fura 2/AM and quin 2/AM into single smooth muscle cells prepared from guinea pig taenia coli. *FEBS Lett*, 220, 89.
26. Erne P, Schachter M, Fabbro D, Miles CM, Sever PS. (1987) Calcium transients in human platelets monitored by aequorin, fura-2 and quin-2: effects of protein kinase C activation and inhibition. *Biochem Biophys Res Commun*, 145, 66.
27. Bibb PC, Cochrane DE, Morel-Laurens N. (1986) Loss of quin 2 accompanies degranulation of mast cells. *FEBS Lett*, 209, 169.
28. James-Kracke MR. (1986) Measurement of cytoplasmic free Ca^{2+} concentration in cultured muscle cells by aequorin and quin 2. *Am J Physiol*, 251, C512.
29. Suko J, Wyskovsky W, Pidlich J, Hauptner R, Plank B, Hellmann G. (1986) Calcium release from calmodulin and its C-terminal or N-terminal halves in the presence of the calmodulin antagonists phenoxybenzamine and melittin measured by stopped-flow fluorescence with Quin 2 and intrinsic tyrosine. Inhibition of calmodulin-dependent protein kinase of cardiac sarcoplasmic reticulum. *Eur J Biochem*, 159, 425.
30. Goligorsky MS, Hruska KA, Loftus DJ, Elson EL. (1986) Alpha 1-adrenergic stimulation and cytoplasmic free calcium concentration in cultured renal proximal tubular cells: evidence for compartmentalization of quin-2 and fura-2. *J Cell Physiol*, 128, 466.
31. Hart GR, Ray KP, Wallis M. (1986) Use of quin 2 to measure calcium concentrations in ovine anterior pituitary cells and the effects of quin 2 on secretion of growth hormone and prolactin. *FEBS Lett*, 203, 77.
32. Williams DA, Fay FS. (1986) Calcium transients and resting levels in isolated smooth muscle cells as monitored with quin 2. *Am J Physiol*, 250, C779.
33. Kao LS, Schneider AS. (1986) Calcium mobilization and catecholamine secretion in adrenal chromaffin cells. A Quin-2 fluorescence study. *J Biol Chem*, 261, 4881.
34. DeFeo TT, Morgan KG. (1986) A comparison of two different indicators: quin 2 and aequorin in isolated single cells and intact strips of ferret portal vein. *Pflugers Arch*, 406, 427.
35. Simon MF, Chap H, Douste-Blazy L. (1986) Selective inhibition of human platelet phospholipase A2 by buffering cytoplasmic calcium with the fluorescent indicator quin 2. Evidence for different calcium sensitivities of phospholipases A2 and C. *Biochim Biophys Acta*, 875, 157.
36. Matsuno K, Katabami F, Koyama M, Abe K, Kobayashi M, Sakurada K, Miyazaki T. (1986) [On the measurement of cytosolic free calcium ion concentration using a fluorescent probe, quin 2]. *Rinsho Ketsueki*, 27, 165.
37. Rao GH, Peller JD, Semba CP, White JG. (1986) Influence of the calcium-sensitive fluorophore, Quin 2, on platelet function. *Blood*, 67, 354.
38. Ohki K, Nagaoka S, Sogami M, Nozawa Y. (1986) Ca^{2+} -translocation activities of phosphatidylinositol, diacylglycerol and phosphatidic acid inferred by quin-2 in artificial membrane systems. *Chem Phys Lipids*, 39, 237.
39. Deber CM, Hsu LC. (1986) Calcium transport by ionophorous peptides in dog and human lymphocytes detected by quin-2 fluorescence. *Biochem Biophys Res Commun*, 134, 731.
40. Utsumi K, Nobori K, Terada S, Miyahara M, Utsumi T. (1985) Continuous fluorometric assay of Ca^{2+} transport by liposomes with Quin 2 entrapped: effect of phospholipase A2 and unsaturated long-chain fatty acids. *Cell Struct Funct*, 10, 339.

41. Sistare FD, Picking RA, Haynes RC, Jr. (1985) Sensitivity of the response of cytosolic calcium in Quin-2-loaded rat hepatocytes to glucagon, adenine nucleosides, and adenine nucleotides. *J Biol Chem*, 260, 12744.
42. Martin SR, Andersson Teleman A, Bayley PM, Drakenberg T, Forsen S. (1985) Kinetics of calcium dissociation from calmodulin and its tryptic fragments. A stopped-flow fluorescence study using Quin 2 reveals a two-domain structure. *Eur J Biochem*, 151, 543.
43. Allen TJ, Baker PF. (1985) Intracellular Ca indicator Quin-2 inhibits Ca²⁺ inflow via Na/Ca exchange in squid axon. *Nature*, 315, 755.
44. Hatayama K, Kambayashi J, Nakamura K, Ohshiro T, Mori T. (1985) Fluorescent Ca²⁺-indicator quin 2 as an intracellular Ca²⁺ antagonist in platelet reaction. *Thromb Res*, 38, 505.
45. Harvey DJ, Godber JF, Timmerman MP, Castell LM, Ashley CC. (1985) Measurement of free Ca²⁺ changes and total Ca²⁺ release in a single striated muscle fibre using the fluorescent indicator quin 2. *Biochem Biophys Res Commun*, 128, 1180.
46. Powers RE, Johnson PC, Houlihan MJ, Saluja AK, Steer ML. (1985) Intracellular Ca²⁺ levels and amylase secretion in Quin 2-loaded mouse pancreatic acini. *Am J Physiol*, 248, C535.
47. Lokesh BR, Kinsella JE. (1985) Intracellular calcium does not appear to be essential for arachidonic acid release from stimulated macrophages as shown by studies with Quin-2. *Biochim Biophys Acta*, 845, 101.
48. Higashihara M, Kume S. (1985) [Calcium-platelet endogenous protein phosphorylation, Quin 2 fluorescence, CTC fluorescence]. *Rinsho Byori, Spec No 62*, 105.
49. Bryant DT. (1985) Quin 2: the dissociation constants of its Ca²⁺ and Mg²⁺ complexes and its use in a fluorimetric method for determining the dissociation of Ca²⁺-protein complexes. *Biochem J*, 226, 613.
50. Johnson PC, Ware JA, Cliveden PB, Smith M, Dvorak AM, Salzman EW. (1985) Measurement of ionized calcium in blood platelets with the photoprotein aequorin. Comparison with Quin 2. *J Biol Chem*, 260, 2069.
51. Uchida MK, Mita M. (1985) Effects of Quin-2, an intracellular Ca chelator, and TMB-8, an intracellular Ca antagonist, on the short-term desensitization to histamine in guinea pig taenia caecum. *Gen Pharmacol*, 16, 449.
52. Salzman EW, Johnson PC, Ware JA. (1985) Measurement of intracellular platelet calcium with aequorin and Quin 2. *Adv Exp Med Biol*, 192, 163.
53. Sheu SS, Sharma VK, Banerjee SP. (1984) Measurement of cytosolic free calcium concentration in isolated rat ventricular myocytes with quin 2. *Circ Res*, 55, 830.
54. Quast U, Labhardt AM, Doyle VM. (1984) Stopped-flow kinetics of the interaction of the fluorescent calcium indicator Quin 2 with calcium ions. *Biochem Biophys Res Commun*, 123, 604.
55. Hirata M, Hashimoto T, Hamachi T, Koga T. (1984) Changes of intracellular free Ca²⁺ in macrophages following N-formyl chemotactic peptide stimulation. Direct measurement by the loading of quin 2. *J Biochem (Tokyo)*, 96, 9.
56. White JR, Ishizaka T, Ishizaka K, Sha'afi R. (1984) Direct demonstration of increased intracellular concentration of free calcium as measured by quin-2 in stimulated rat peritoneal mast cell. *Proc Natl Acad Sci U S A*, 81, 3978.
57. Bayley P, Ahlstrom P, Martin SR, Forsen S. (1984) The kinetics of calcium binding to calmodulin: Quin 2 and ANS stopped-flow fluorescence studies. *Biochem Biophys Res Commun*, 120, 185.
58. Ashley RH, Brammer MJ, Marchbanks R. (1984) Measurement of intrasynaptosomal free calcium by using the fluorescent indicator quin-2. *Biochem J*, 219, 149.
59. Muallem S, Blissard D, Sachs G. (1984) Single cell recording of quin 2 fluorescence. *Prog Clin Biol Res*, 168, 151.
60. Rogers J, Hesketh TR, Smith GA, Beaven MA, Metcalfe JC, Johnson P, Garland PB. (1983) Intracellular pH and free calcium changes in single cells using quene 1 and quin 2 probes and fluorescence microscopy. *FEBS Lett*, 161, 21.

61. Charest R, Blackmore PF, Berthon B, Exton JH. (1983) Changes in free cytosolic Ca²⁺ in hepatocytes following alpha 1-adrenergic stimulation. Studies on Quin-2-loaded hepatocytes. *J Biol Chem*, 258, 8769.