

References for Products 21416 and 21417

1. Strathmann M, Wingender J. (2004) Use of an oxonol dye in combination with confocal laser scanning microscopy to monitor damage to *Staphylococcus aureus* cells during colonisation of silver-coated vascular grafts. *Int J Antimicrob Agents*, 24, 234.
2. Portele A, Lenz J, Hofer M. (1997) Estimation of membrane potential $\Delta\psi$ in reconstituted plasma membrane vesicles using a numerical model of oxonol VI distribution. *J Bioenerg Biomembr*, 29, 603.
3. Jepras RI, Paul FE, Pearson SC, Wilkinson MJ. (1997) Rapid assessment of antibiotic effects on *Escherichia coli* by bis-(1,3-dibutylbarbituric acid) trimethine oxonol and flow cytometry. *Antimicrob Agents Chemother*, 41, 2001.
4. Dall'Asta V, Gatti R, Orlandini G, Rossi PA, Rotoli BM, Sala R, Bussolati O, Gazzola GC. (1997) Membrane potential changes visualized in complete growth media through confocal laser scanning microscopy of bis-oxonol-loaded cells. *Exp Cell Res*, 231, 260.
5. Kukkonen JP, Hautala R, Akerman KE. (1996) Muscarinic depolarization of SH-SY5Y human neuroblastoma cells as determined using oxonol V. *Neurosci Lett*, 212, 57.
6. Knauf PA, Law FY, Hahn K. (1995) An oxonol dye is the most potent known inhibitor of band 3-mediated anion exchange. *Am J Physiol*, 269, C1073.
7. Arreola J, Hallows KR, Knauf PA. (1995) Volume-activated chloride channels in HL-60 cells: potent inhibition by an oxonol dye. *Am J Physiol*, 269, C1063.
8. Lopez-Amoros R, Comas J, Vives-Rego J. (1995) Flow cytometric assessment of *Escherichia coli* and *Salmonella typhimurium* starvation-survival in seawater using rhodamine 123, propidium iodide, and oxonol. *Appl Environ Microbiol*, 61, 2521.
9. Ahmed I, Krishnamoorthy G. (1994) Anomalous response of oxonol-V to membrane potential in mitochondrial proton pumps. *Biochim Biophys Acta*, 1188, 131.
10. Epps DE, Wolfe ML, Groppi V. (1994) Characterization of the steady-state and dynamic fluorescence properties of the potential-sensitive dye bis-(1,3-dibutylbarbituric acid)trimethine oxonol (Dibac4(3)) in model systems and cells. *Chem Phys Lipids*, 69, 137.
11. Gamalei IA, Kaulin AB, Kirpichnikova KM. (1991) [The measurement of the macrophage membrane potential by using an oxonol fluorescent probe]. *Tsitologija*, 33, 60.
12. Kitayama S, Ohtsuki H, Morita K, Dohi T, Tsujimoto A. (1990) Bis-oxonol experiment on plasma membrane potentials of bovine adrenal chromaffin cells: depolarizing stimuli and their possible interaction. *Neurosci Lett*, 116, 275.
13. Cooper CE, Bruce D, Nicholls P. (1990) Use of oxonol V as a probe of membrane potential in proteoliposomes containing cytochrome oxidase in the submitochondrial orientation. *Biochemistry*, 29, 3859.
14. Clarke RJ, Apell HJ. (1989) A stopped-flow kinetic study of the interaction of potential-sensitive oxonol dyes with lipid vesicles. *Biophys Chem*, 34, 225.
15. George EB, Nyirjesy P, Pratap PR, Freedman JC, Waggoner AS. (1988) Impermeant potential-sensitive oxonol dyes: III. The dependence of the absorption signal on membrane potential. *J Membr Biol*, 105, 55.
16. Nyirjesy P, George EB, Gupta RK, Basson M, Pratap PR, Freedman JC, Raman K, Waggoner AS. (1988) Impermeant potential-sensitive oxonol dyes: II. The dependence of the absorption signal on the length of alkyl substituents attached to the dye. *J Membr Biol*, 105, 45.
17. George EB, Nyirjesy P, Basson M, Ernst LA, Pratap PR, Freedman JC, Waggoner AS. (1988) Impermeant potential-sensitive oxonol dyes: I. Evidence for an "on-off" mechanism. *J Membr Biol*, 103, 245.
18. Apell HJ, Bersch B. (1987) Oxonol VI as an optical indicator for membrane potentials in lipid vesicles. *Biochim Biophys Acta*, 903, 480.
19. Mohr FC, Fewtrell C. (1987) IgE receptor-mediated depolarization of rat basophilic leukemia cells measured with the fluorescent probe bis-oxonol. *J Immunol*, 138, 1564.

20. Wilson HA, Chused TM. (1985) Lymphocyte membrane potential and Ca²⁺-sensitive potassium channels described by oxonol dye fluorescence measurements. *J Cell Physiol*, 125, 72.
21. Bashford CL, Alder GM, Gray MA, Micklem KJ, Taylor CC, Turek PJ, Pasternak CA. (1985) Oxonol dyes as monitors of membrane potential: the effect of viruses and toxins on the plasma membrane potential of animal cells in monolayer culture and in suspension. *J Cell Physiol*, 123, 326.
22. van Walraven HS, Krab K, Hagendoorn MJ, Kraayenhof R. (1985) The use of carotenoids and oxonol VI as probes for membrane potential in proteoliposomes. *FEBS Lett*, 184, 96.
23. Pringle MJ, Sanadi DR. (1984) Effects of Cd²⁺ on ATP-driven membrane potential in beef heart mitochondrial H⁺-ATPase: a study using the voltage-sensitive probe oxonol VI. *Membr Biochem*, 5, 225.
24. Freedman JC, Novak TS. (1983) Membrane potentials associated with Ca-induced K conductance in human red blood cells: studies with a fluorescent oxonol dye, WW 781. *J Membr Biol*, 72, 59.
25. Smith JC, Hallidy L, Topp MR. (1981) The behavior of the fluorescence lifetime and polarization of oxonol potential-sensitive extrinsic probes in solution and in beef heart submitochondrial particles. *J Membr Biol*, 60, 173.
26. Scherman D, Henry JP. (1980) Oxonol-V as a probe of chromaffin granule membrane potentials. *Biochim Biophys Acta*, 599, 150.
27. Russell JT, Beeler T, Martonosi A. (1979) Optical probe responses on sarcoplasmic reticulum. Merocyanine and oxonol dyes. *J Biol Chem*, 254, 2047.
28. Bashford CL, Chance B, Prince RC. (1979) Oxonol dyes as monitors of membrane potential. Their behavior in photosynthetic bacteria. *Biochim Biophys Acta*, 545, 46.
29. Smith JC, Chance B. (1979) Kinetics of the potential-sensitive extrinsic probe oxonol VI in beef heart submitochondrial particles. *J Membr Biol*, 46, 255.