

References for Product 21492

1. Saldanha C, Santos NC, Martins-Silva J. (2002) Fluorescent probes DPH, TMA-DPH and C17-HC induce erythrocyte exovesiculation. *J Membr Biol*, 190, 75.
2. Matko J, Nagy P. (1997) Fluorescent lipid probes 12-AS and TMA-DPH report on selective, purinergically induced fluidity changes in plasma membranes of lymphoid cells. *J Photochem Photobiol B*, 40, 120.
3. Simonetti O, Ferretti G, Offidani AM, Gervasi P, Curatola G, Bossi G. (1996) Plasma membrane fluidity of keratinocytes of normal and psoriatic skin: a study using fluorescence anisotropy of trimethylammoniumdiphenylhexatriene (TMA-DPH). *Arch Dermatol Res*, 288, 51.
4. Illinger D, Duportail G, Mely Y, Poirel-Morales N, Gerard D, Kuhry JG. (1995) A comparison of the fluorescence properties of TMA-DPH as a probe for plasma membrane and for endocytic membrane. *Biochim Biophys Acta*, 1239, 58.
5. Obsil T, Amler E, Kotyk A, Pavlicek Z. (1995) Enzymically generated electron-excited species increase the membrane order parameter without changing the rotational relaxation time of TMA-DPH. *Gen Physiol Biophys*, 14, 393.
6. Illinger D, Kuhry JG. (1994) The kinetic aspects of intracellular fluorescence labeling with TMA-DPH support the maturation model for endocytosis in L929 cells. *J Cell Biol*, 125, 783.
7. Herman P, Konopasek I, Plasek J, Svobodova J. (1994) Time-resolved polarized fluorescence studies of the temperature adaptation in *Bacillus subtilis* using DPH and TMA-DPH fluorescent probes. *Biochim Biophys Acta*, 1190, 1.
8. Ferretti G, Zolese G, Curatola G, Jezequel AM, Benedetti A. (1993) Membrane heterogeneity in isolated rat hepatocytes and liver plasma membrane subfractions: a comparative study using DPH and its cationic derivative TMA-DPH. *Biochim Biophys Acta*, 1147, 245.
9. Ferretti G, Tangorra A, Zolese G, Curatola G. (1993) Properties of a phosphatidylcholine derivative of diphenyl hexatriene (DPH-PC) in lymphocyte membranes. A comparison with DPH and the cationic derivative TMA-DPH using static and dynamic fluorescence. *Membr Biochem*, 10, 17.
10. Illinger D, Italiano L, Beck JP, Waltzinger C, Kuhry JG. (1993) Comparative evolution of endocytosis levels and of the cell surface area during the L929 cell cycle: a fluorescence study with TMA-DPH. *Biol Cell*, 79, 265.
11. Maire C, Bouchy M, Donner M, Andre JC. (1992) Membrane labelling by fluorescent probes: incorporation of TMA-DPH in erythrocyte membranes. *Biorheology*, 29, 507.
12. Toplak H, Batchiulis V, Hermetter A, Hunziker T, Honegger UE, Wiesmann UN. (1990) Effects of culture and incubation conditions on membrane fluidity in monolayers of cultured cells measured as fluorescence anisotropy using trimethylammoniumdiphenylhexatriene (TMA-DPH). *Biochim Biophys Acta*, 1028, 67.
13. Zolese G, Gratton E, Curatola G. (1990) Phosphatidic acid affects structural organization of phosphatidylcholine liposomes. A study of 1,6-diphenyl-1,3,5-hexatriene (DPH) and 1-(4-trimethylammonium-phenyl)-6-phenyl-1,3,5-hexatriene (TMA-DPH) fluorescence decay using distributional analysis. *Chem Phys Lipids*, 55, 29.
14. Benedetti A, Ferretti G, Curatola G, Brunelli E, Jezequel AM, Orlandi F. (1989) Plasma membrane fluidity in isolated rat hepatocytes: comparative study using DPH and TMA-DPH as fluorescent probes. *J Gastroenterol Hepatol*, 4, 221.
15. Illinger D, Kubina M, Duportail G, Poindron P, Bartholeyns J, Kuhry JG. (1989) TMA-DPH a fluorescent probe of membrane dynamics in living cells. How to use it in phagocytosis. *Cell Biophys*, 14, 17.
16. Hitzemann RJ, Harris RA. (1984) Developmental changes in synaptic membrane fluidity: a comparison of 1,6-diphenyl-1,3,5-hexatriene (DPH) and 1-[4-(trimethylamino)phenyl]-6-phenyl-1,3,5-hexatriene (TMA-DPH). *Brain Res*, 316, 113.

17. Kuhry JG, Fonteneau P, Duportail G, Maechling C, Laustriat G. (1983) TMA-DPH: a suitable fluorescence polarization probe for specific plasma membrane fluidity studies in intact living cells. *Cell Biophys*, 5, 129.