

References for Product 22205

1. Wiosetek-Reske AM, Wysocki S. (2006) Spectral studies of N-nonyl acridine orange in anionic, cationic and neutral surfactants. *Spectrochim Acta A Mol Biomol Spectrosc*, 64, 1118.
2. Gohil VM, Gvozdenovic-Jeremic J, Schlame M, Greenberg ML. (2005) Binding of 10-N-nonyl acridine orange to cardiolipin-deficient yeast cells: implications for assay of cardiolipin. *Anal Biochem*, 343, 350.
3. Wiosetek-Reske AM, Wysocki S, Bak GW. (2005) Determination of dipole moment in the ground and excited state by experimental and theoretical methods of N-nonyl acridine orange. *Spectrochim Acta A Mol Biomol Spectrosc*, 62, 1172.
4. Garcia Fernandez MI, Ceccarelli D, Muscatello U. (2004) Use of the fluorescent dye 10-N-nonyl acridine orange in quantitative and location assays of cardiolipin: a study on different experimental models. *Anal Biochem*, 328, 174.
5. Jacobson J, Duchen MR, Heales SJ. (2002) Intracellular distribution of the fluorescent dye nonyl acridine orange responds to the mitochondrial membrane potential: implications for assays of cardiolipin and mitochondrial mass. *J Neurochem*, 82, 224.
6. Mileykovskaya E, Dowhan W, Birke RL, Zheng D, Lutterodt L, Haines TH. (2001) Cardiolipin binds nonyl acridine orange by aggregating the dye at exposed hydrophobic domains on bilayer surfaces. *FEBS Lett*, 507, 187.
7. Keij JF, Bell-Prince C, Steinkamp JA. (2000) Staining of mitochondrial membranes with 10-nonyl acridine orange, MitoFluor Green, and MitoTracker Green is affected by mitochondrial membrane potential altering drugs. *Cytometry*, 39, 203.
8. Mileykovskaya E, Dowhan W. (2000) Visualization of phospholipid domains in *Escherichia coli* by using the cardiolipin-specific fluorescent dye 10-N-nonyl acridine orange. *J Bacteriol*, 182, 1172.
9. Gallet PF, Maftah A, Petit JM, Denis-Gay M, Julien R. (1995) Direct cardiolipin assay in yeast using the red fluorescence emission of 10-N-nonyl acridine orange. *Eur J Biochem*, 228, 113.
10. Petit JM, Maftah A, Ratinaud MH, Julien R. (1992) 10N-nonyl acridine orange interacts with cardiolipin and allows the quantification of this phospholipid in isolated mitochondria. *Eur J Biochem*, 209, 267.
11. Lizard G, Chardonnet Y, Chignol MC, Thivolet J. (1990) Evaluation of mitochondrial content and activity with nonyl-acridine orange and rhodamine 123: flow cytometric analysis and comparison with quantitative morphometry. Comparative analysis by flow cytometry and quantitative morphometry of mitochondrial content and activity. *Cytotechnology*, 3, 179.
12. Maftah A, Petit JM, Julien R. (1990) Specific interaction of the new fluorescent dye 10-N-nonyl acridine orange with inner mitochondrial membrane. A lipid-mediated inhibition of oxidative phosphorylation. *FEBS Lett*, 260, 236.
13. Leprat P, Ratinaud MH, Maftah A, Petit JM, Julien R. (1990) Use of nonyl acridine orange and rhodamine 123 to follow biosynthesis and functional assembly of mitochondrial membrane during L1210 cell cycle. *Exp Cell Res*, 186, 130.
14. Maftah A, Petit JM, Ratinaud MH, Julien R. (1989) 10-N nonyl-acridine orange: a fluorescent probe which stains mitochondria independently of their energetic state. *Biochem Biophys Res Commun*, 164, 185.
15. Benel L, Ronot X, Mounolou JC, Gaudemer F, Adolphe M. (1989) Compared flow cytometric analysis of mitochondria using 10-n-nonyl acridine orange and rhodamine 123. *Basic Appl Histochem*, 33, 71.
16. Ratinaud MH, Leprat P, Julien R. (1988) In situ flow cytometric analysis of nonyl acridine orange-stained mitochondria from splenocytes. *Cytometry*, 9, 206.