References for Product 22780

- 1. Bryce SM, Bemis JC, Avlasevich SL, Dertinger SD. (2007) In vitro micronucleus assay scored by flow cytometry provides a comprehensive evaluation of cytogenetic damage and cytotoxicity. Mutat Res, 630, 78.
- 2. Fotakis G, Timbrell JA. (2006) In vitro cytotoxicity assays: comparison of LDH, neutral red, MTT and protein assay in hepatoma cell lines following exposure to cadmium chloride. Toxicol Lett, 160, 171.
- 3. Vichai V, Kirtikara K. (2006) Sulforhodamine B colorimetric assay for cytotoxicity screening. Nat Protoc, 1, 1112.
- 4. Wang X, Ge J, Wang K, Qian J, Zou Y. (2006) Evaluation of MTT assay for measurement of emodin-induced cytotoxicity. Assay Drug Dev Technol, 4, 203.
- Kasatori N, Ishikawa F, Ueyama M, Urayama T. (2005) A differential assay of NK-cellmediated cytotoxicity in K562 cells revealing three sequential membrane impairment steps using three-color flow-cytometry. J Immunol Methods, 307, 41.
- 6. Zimmermann SY, Esser R, Rohrbach E, Klingebiel T, Koehl U. (2005) A novel four-colour flow cytometric assay to determine natural killer cell or T-cell-mediated cellular cytotoxicity against leukaemic cells in peripheral or bone marrow specimens containing greater than 20% of normal cells. J Immunol Methods, 296, 63.
- 7. Ewen C, Kane KP, Shostak I, Griebel PJ, Bertram EM, Watts TH, Bleackley RC, McElhaney JE. (2003) A novel cytotoxicity assay to evaluate antigen-specific CTL responses using a colorimetric substrate for Granzyme B. J Immunol Methods, 276, 89.
- 8. Nouri AM, Mansouri M, Hussain RF, Dos Santos AV, Oliver RT. (1995) Super-sensitive epithelial cell line and colorimetric assay to replace the conventional K562 target and chromium release assay for assessment of non-MHC-restricted cytotoxicity. J Immunol Methods, 180, 63.