

References for Product 23056

1. McLaughlin RW, De Stigter JK, Sikkink LA, Baden EM, Ramirez-Alvarado M. (2006) The effects of sodium sulfate, glycosaminoglycans, and Congo red on the structure, stability, and amyloid formation of an immunoglobulin light-chain protein. *Protein Sci*, 15, 1710.
2. Inestrosa NC, Alvarez A, Dinamarca MC, Perez-Acle T, Colombres M. (2005) Acetylcholinesterase-amyloid-beta-peptide interaction: effect of Congo Red and the role of the Wnt pathway. *Curr Alzheimer Res*, 2, 301.
3. Inouye H, Kirschner DA. (2005) Alzheimer's beta-amyloid: insights into fibril formation and structure from Congo red binding. *Subcell Biochem*, 38, 203.
4. Giorgadze TA, Shiina N, Baloch ZW, Tomaszewski JE, Gupta PK. (2004) Improved detection of amyloid in fat pad aspiration: an evaluation of Congo red stain by fluorescent microscopy. *Diagn Cytopathol*, 31, 300.
5. Sabate R, Estelrich J. (2003) Pinacyanol as effective probe of fibrillar beta-amyloid peptide: comparative study with Congo Red. *Biopolymers*, 72, 455.
6. Kim YS, Randolph TW, Manning MC, Stevens FJ, Carpenter JF. (2003) Congo red populates partially unfolded states of an amyloidogenic protein to enhance aggregation and amyloid fibril formation. *J Biol Chem*, 278, 10842.
7. Rybarska J, Piekarska B, Stopa B, Zemanek G, Konieczny L, Nowak M, Krol M, Roterman I, Szymczakiewicz-Multanowska A. (2001) Evidence that supramolecular Congo red is the sole ligation form of this dye for L chain lambda derived amyloid proteins. *Folia Histochem Cytobiol*, 39, 307.
8. Roterman I, Krul M, Nowak M, Konieczny L, Rybarska J, Stopa B, Piekarska B, Zemanek G. (2001) Why Congo red binding is specific for amyloid proteins - model studies and a computer analysis approach. *Med Sci Monit*, 7, 771.
9. Khurana R, Uversky VN, Nielsen L, Fink AL. (2001) Is Congo red an amyloid-specific dye? *J Biol Chem*, 276, 22715.
10. Michael H, Brandt LJ, Tanaka KE, Berkowitz D, Cardillo M, Weidenheim K. (2001) Congo-red negative colonic amyloid with scalloping of the valvulae conniventes. *Gastrointest Endosc*, 53, 653.
11. Hirakura Y, Yiu WW, Yamamoto A, Kagan BL. (2000) Amyloid peptide channels: blockade by zinc and inhibition by Congo red (amyloid channel block). *Amyloid*, 7, 194.
12. Linke RP. (2000) Highly sensitive diagnosis of amyloid and various amyloid syndromes using Congo red fluorescence. *Virchows Arch*, 436, 439.
13. Chrambach A, Brining SK. (2000) Gel electrophoretic distinction between Congo Red nonreactive beta-amyloid (1-42) and beta-amyloid (1-40). *Electrophoresis*, 21, 760.
14. Hirakura Y, Lin MC, Kagan B. (1999) Erratum: hirakura Y, lin M-C, kagan BL. 1999. Alzheimer amyloid Abeta1-42 channels: effects of solvent, pH, and congo red. *J neurosci res* 57:458-466. *J Neurosci Res*, 58, 726.
15. Klunk WE, Jacob RF, Mason RP. (1999) Quantifying amyloid by congo red spectral shift assay. *Methods Enzymol*, 309, 285.
16. Hirakura Y, Lin MC, Kagan BL. (1999) Alzheimer amyloid abeta1-42 channels: effects of solvent, pH, and Congo Red. *J Neurosci Res*, 57, 458.
17. Klunk WE, Jacob RF, Mason RP. (1999) Quantifying amyloid beta-peptide (Abeta) aggregation using the Congo red-Abeta (CR-abeta) spectrophotometric assay. *Anal Biochem*, 266, 66.
18. Carter DB, Chou KC. (1998) A model for structure-dependent binding of Congo red to Alzheimer beta-amyloid fibrils. *Neurobiol Aging*, 19, 37.
19. Podlisny MB, Walsh DM, Amarante P, Ostaszewski BL, Stimson ER, Maggio JE, Teplow DB, Selkoe DJ. (1998) Oligomerization of endogenous and synthetic amyloid beta-protein at nanomolar levels in cell culture and stabilization of monomer by Congo red. *Biochemistry*, 37, 3602.
20. Gupta-Bansal R, Brunden KR. (1998) Congo red inhibits proteoglycan and serum amyloid P binding to amyloid beta fibrils. *J Neurochem*, 70, 292.

21. Abe K, Kato M, Saito H. (1997) Congo red reverses amyloid beta protein-induced cellular stress in astrocytes. *Neurosci Res*, 29, 129.
22. Ashburn TT, Han H, McGuinness BF, Lansbury PT, Jr. (1996) Amyloid probes based on Congo Red distinguish between fibrils comprising different peptides. *Chem Biol*, 3, 351.
23. Linke RP, Gartner HV, Michels H. (1995) High-sensitivity diagnosis of AA amyloidosis using Congo red and immunohistochemistry detects missed amyloid deposits. *J Histochem Cytochem*, 43, 863.
24. Burgevin MC, Passat M, Daniel N, Capet M, Doble A. (1994) Congo red protects against toxicity of beta-amyloid peptides on rat hippocampal neurones. *Neuroreport*, 5, 2429.
25. Lorenzo A, Yankner BA. (1994) Beta-amyloid neurotoxicity requires fibril formation and is inhibited by congo red. *Proc Natl Acad Sci U S A*, 91, 12243.
26. Askanas V, Engel WK, Alvarez RB. (1993) Enhanced detection of congo-red-positive amyloid deposits in muscle fibers of inclusion body myositis and brain of Alzheimer's disease using fluorescence technique. *Neurology*, 43, 1265.
27. Turnell WG, Finch JT. (1992) Binding of the dye congo red to the amyloid protein pig insulin reveals a novel homology amongst amyloid-forming peptide sequences. *J Mol Biol*, 227, 1205.
28. Yang GC, Nieto R, Stachura I, Gallo GR. (1992) Ultrastructural immunohistochemical localization of polyclonal IgG, C3, and amyloid P component on the congo red-negative amyloid-like fibrils of fibrillary glomerulopathy. *Am J Pathol*, 141, 409.
29. Klunk WE, Pettegrew JW, Abraham DJ. (1989) Quantitative evaluation of congo red binding to amyloid-like proteins with a beta-pleated sheet conformation. *J Histochem Cytochem*, 37, 1273.
30. Egner E. (1981) [Light microscopic amyloid diagnosis with Congo red and phenol reaction]. *Pathologie*, 2, 233.
31. Linder E, Lehto VP, Virtanen I. (1979) Amyloid-like green birefringence in cytoskeletal 10 nm filaments after staining with Congo red. *Acta Pathol Microbiol Scand [A]*, 87A, 299.
32. Meloan SN, Puchtler H. (1978) Demonstration of amyloid with Mesitol WLS-Congo Red: application of a textile auxiliary to histochemistry. *Histochemistry*, 58, 163.
33. Stokes G. (1976) An improved Congo red method for amyloid. *Med Lab Sci*, 33, 79.
34. Brigger D, Muckle RJ. (1975) Comparison of Sirius red and Congo red as stains for amyloid in animal tissues. *J Histochem Cytochem*, 23, 84.
35. Taylor DL, Allen RD, Benditt EP. (1974) Determination of the polarization optical properties of the amyloid-Congo red complex by phase modulation microspectrophotometry. *J Histochem Cytochem*, 22, 1105.
36. Cooper JH. (1974) Selective amyloid staining as a function of amyloid composition and structure. Histochemical analysis of the alkaline Congo red, standardized toluidine blue, and iodine methods. *Lab Invest*, 31, 232.
37. Glenner GG, Eanes ED, Page DL. (1972) The relation of the properties of Congo red-stained amyloid fibrils to the -conformation. *J Histochem Cytochem*, 20, 821.
38. Ruinen L, Smit-van Dijk W, Pros M, Scholten JH, Mandema E. (1972) The binding of Congo red and toluidine blue to isolated amyloid. *Acta Pathol Microbiol Scand [A]*, 233, 122.
39. Katenkamp D, Stiller D. (1972) [Polarization microscopic and histochemical studies on the Congo red-staining of amyloid]. *Histochemie*, 29, 37.
40. Eastwood H, Cole KR. (1971) Staining of amyloid by buffered Congo red in 50 per cent ethanol. *Stain Technol*, 46, 208.
41. Romhanyi G. (1971) Selective differentiation between amyloid and connective tissue structures based on the collagen specific topo-optical staining reaction with congo red. *Virchows Arch A Pathol Pathol Anat*, 354, 209.
42. Benditt EP, Eriksen N, Berglund C. (1970) Congo red dichroism with dispersed amyloid fibrils, an extrinsic cotton effect. *Proc Natl Acad Sci U S A*, 66, 1044.
43. Soppitt GD, Pennock CA. (1969) Interaction of water-soluble amyloid fibrils with Congo Red and thioflavine T. *Clin Chim Acta*, 26, 165.
44. Reissenweber NJ, Decaro J. (1969) Dichroism with congo red. A specific test for amyloid? *Virchows Arch A Pathol Pathol Anat*, 347, 254.

45. Ashkenazi Y, Hersko C, Gafni J, Sohar E, Heller H. (1967) Chemical aspects of amyloid-Congo red binding. *Isr J Med Sci*, 3, 572.
46. Kelenyi G. (1967) Thioflavin S fluorescent and Congo red anisotropic stainings in the histologic demonstration of amyloid. *Acta Neuropathol (Berl)*, 7, 336.
47. Hewlett BR. (1966) Studies on the staining of amyloid with Congo red. *Can J Med Technol*, 28, 248.
48. Shirahama T, Cohen AS. (1966) A congo red staining method for epoxy-embedded amyloid. *J Histochem Cytochem*, 14, 725.
49. Puchtler H, Sweat F. (1965) Congo red as a stain for fluorescence microscopy of amyloid. *J Histochem Cytochem*, 13, 693.
50. Wolman M, Bubis JJ. (1965) The cause of the green polarization color of amyloid stained with Congo red. *Histochemie*, 4, 351.
51. Jamrozik T, Michalowicz R, Zukowska W. (1962) [Evaluation of the Congo red test as a method for early detection of amyloid changes in tuberculous children.]. *Pediatr Pol*, 37, 825.
52. Missmahl HP. (1957) [Polarization microscopic contribution to Congo red staining of amyloid.]. *Z Wiss Mikrosk*, 63, 133.
53. Johansson GA, Pfeiffer HH. (1954) On the amyloid-Congo red complex in histological sections and the genesis of the amyloid substance. *Acta Anat (Basel)*, 20, 285.
54. Lanari A, Patalano A, Soubrie A. (1951) [Rate of Congo red adsorption by amyloid substance in vivo (demonstration by catheterization of the suprahepatic vein).]. *Dtsch Med Wochenschr*, 76, 730.
55. Lanari A, Patalano A, Soubrie A. (1951) [Rate of fixation of Congo red by amyloid substance in vivo; demonstration by catheterization of the suprahepatic vein.]. *Medicina (B Aires)*, 11, 77.
56. Unger PN, Zuckerbrod M, Beck GJ, Steele JM, Porosowska Y. (1948) Study of the Disappearance of Congo Red from the Blood of Non-Amyloid Subjects and Patients with Amyloidosis. *J Clin Invest*, 27, 111.