

References for Product 36701

1. Mita S, Suzuki H, Akita H, Hayashi H, Onuki R, Hofmann AF, Sugiyama Y. (2006) Inhibition of bile acid transport across Na⁺/taurocholate cotransporting polypeptide (SLC10A1) and bile salt export pump (ABCB 11)-coexpressing LLC-PK1 cells by cholestasis-inducing drugs. *Drug Metab Dispos*, 34, 1575.
2. Perez LM, Milkiewicz P, Elias E, Coleman R, Sanchez Pozzi EJ, Roma MG. (2006) Oxidative stress induces internalization of the bile salt export pump, Bsep, and bile salt secretory failure in isolated rat hepatocyte couplets: a role for protein kinase C and prevention by protein kinase A. *Toxicol Sci*, 91, 150.
3. Wang W, Soroka CJ, Mennone A, Rahner C, Harry K, Pypaert M, Boyer JL. (2006) Radixin is required to maintain apical canalicular membrane structure and function in rat hepatocytes. *Gastroenterology*, 131, 878.
4. Sidler Pfandler MA, Hochli M, Inderbitzin D, Meier PJ, Stieger B. (2004) Small hepatocytes in culture develop polarized transporter expression and differentiation. *J Cell Sci*, 117, 4077.
5. Beuers U, Denk GU, Soroka CJ, Wimmer R, Rust C, Paumgartner G, Boyer JL. (2003) Tauroolithocholic acid exerts cholestatic effects via phosphatidylinositol 3-kinase-dependent mechanisms in perfused rat livers and rat hepatocyte couplets. *J Biol Chem*, 278, 17810.
6. Kida M, Mano Y, Ueno Y, Kobayashi K, Goto J, Ishii M, Shimosegawa T. (2003) Vectorial transport of bile acids in immortalized mouse bile duct cells. *Hepatology*, 37, 151.
7. Roman ID, Fernandez-Moreno MD, Fueyo JA, Roma MG, Coleman R. (2003) Cyclosporin A induced internalization of the bile salt export pump in isolated rat hepatocyte couplets. *Toxicol Sci*, 71, 276.
8. Milkiewicz P, Roma MG, Elias E, Coleman R. (2002) Hepatoprotection with tauroursodeoxycholate and beta muricholate against tauroolithocholate induced cholestasis: involvement of signal transduction pathways. *Gut*, 51, 113.
9. Crocenzi FA, Sanchez Pozzi EJ, Pellegrino JM, Favre CO, Rodriguez Garay EA, Mottino AD, Coleman R, Roma MG. (2001) Beneficial effects of silymarin on estrogen-induced cholestasis in the rat: a study in vivo and in isolated hepatocyte couplets. *Hepatology*, 34, 329.
10. Holzinger F, Krahenbuhl L, Schteingart CD, Ton-Nu HT, Hofmann AF. (2001) Use of a fluorescent bile acid to enhance visualization of the biliary tract and bile leaks during laparoscopic surgery in rabbits. *Surg Endosc*, 15, 209.
11. Milkiewicz P, Mills CO, Hubscher SG, Cardenas R, Cardenas T, Williams A, Elias E. (2001) Visualization of the transport of primary and secondary bile acids across liver tissue in rats: in vivo study with fluorescent bile acids. *J Hepatology*, 34, 4.
12. Wustner D, Mukherjee S, Maxfield FR, Muller P, Herrmann A. (2001) Vesicular and nonvesicular transport of phosphatidylcholine in polarized HepG2 cells. *Traffic*, 2, 277.
13. Cantz T, Nies AT, Brom M, Hofmann AF, Keppler D. (2000) MRP2, a human conjugate export pump, is present and transports fluo 3 into apical vacuoles of Hep G2 cells. *Am J Physiol Gastrointest Liver Physiol*, 278, G522.
14. Coleman R, Roma MG. (2000) Hepatocyte couplets. *Biochem Soc Trans*, 28, 136.
15. Michael S, Thole M, Dillmann R, Fahr A, Drewe J, Fricker G. (2000) Improvement of intestinal peptide absorption by a synthetic bile acid derivative, cholylsarcosine. *Eur J Pharm Sci*, 10, 133.
16. Milkiewicz P, Saksena S, Cardenas T, Mills CO, Elias E. (2000) Plasma elimination of cholyl-lysyl-fluorescein (CLF): a pilot study in patients with liver cirrhosis. *Liver*, 20, 330.
17. Roma MG, Milkiewicz P, Elias E, Coleman R. (2000) Control by signaling modulators of the sorting of canalicular transporters in rat hepatocyte couplets: role of the cytoskeleton. *Hepatology*, 32, 1342.
18. Mills CO, Milkiewicz P, Muller M, Roma MG, Havinga R, Coleman R, Kuipers F, Jansen PL, Elias E. (1999) Different pathways of canalicular secretion of sulfated and non-

- sulfated fluorescent bile acids: a study in isolated hepatocyte couplets and TR- rats. *J Hepatol*, 31, 678.
19. Orsler DJ, Ahmed-Choudhury J, Chipman JK, Hammond T, Coleman R. (1999) ANIT-induced disruption of biliary function in rat hepatocyte couplets. *Toxicol Sci*, 47, 203.
 20. Roma MG, Ahmed-Choudhury J, Coleman R. (1999) The protein kinase inhibitor 1-(5-isoquinolinylsulfonyl)-2-methyl-piperzine (H-7) prevents and reverses Ca(2+)-mediated injury in isolated rat hepatocyte couplets. *Toxicol Appl Pharmacol*, 161, 192.
 21. Ahmed-Choudhury J, Orsler DJ, Coleman R. (1998) Hepatobiliary effects of tertiary-butylhydroperoxide (tBOOH) in isolated rat hepatocyte couplets. *Toxicol Appl Pharmacol*, 152, 270.
 22. Bravo P, Bender V, Cassio D. (1998) Efficient in vitro vectorial transport of a fluorescent conjugated bile acid analogue by polarized hepatic hybrid WIF-B and WIF-B9 cells. *Hepatology*, 27, 576.
 23. Briz O, el-Mir MY, Bravo P, Villanueva GR, Marin JJ. (1998) Fetal excretion of the fluorescent bile acid derivative cholyglycylamido-fluorescein (FITC-GC) by the rat placenta-maternal liver tandem. *Placenta*, 19, 119.
 24. El-Mir MY, Serrano MA, Macias RI, Dominguez MF, Monte MJ, Marin JJ. (1998) In vitro test to determine the effect of cytostatic drugs on co-cultured rat hepatocytes and hepatoma cells. *Int J Exp Pathol*, 79, 109.
 25. Holzinger F, Schteingart CD, Ton-Nu HT, Cerre C, Steinbach JH, Yeh HZ, Hofmann AF. (1998) Transport of fluorescent bile acids by the isolated perfused rat liver: kinetics, sequestration, and mobilization. *Hepatology*, 28, 510.
 26. Roma MG, Stone V, Shaw R, Coleman R. (1998) Vasopressin-induced disruption of actin cytoskeletal organization and canalicular function in isolated rat hepatocyte couplets: possible involvement of protein kinase C. *Hepatology*, 28, 1031.
 27. El-Seaidy AZ, Mills CO, Elias E, Crawford JM. (1997) Lack of evidence for vesicle trafficking of fluorescent bile salts in rat hepatocyte couplets. *Am J Physiol*, 272, G298.
 28. Holzinger F, Schteingart CD, Ton-Nu HT, Eming SA, Monte MJ, Hagey LR, Hofmann AF. (1997) Fluorescent bile acid derivatives: relationship between chemical structure and hepatic and intestinal transport in the rat. *Hepatology*, 26, 1263.
 29. Milkiewicz P, Baiocchi L, Mills CO, Ahmed M, Khalaf H, Keogh A, Baker J, Elias E. (1997) Plasma clearance of cholylyl-fluorescein: a pilot study in humans. *J Hepatol*, 27, 1106.
 30. Mills CO, Milkiewicz P, Saraswat V, Elias E. (1997) Cholylyl-fluorescein and related lysyl-fluorescein conjugated bile acid analogues. *Yale J Biol Med*, 70, 447.
 31. Roma MG, Orsler DJ, Coleman R. (1997) Canalicular retention as an in vitro assay of tight junctional permeability in isolated hepatocyte couplets: effects of protein kinase modulation and cholestatic agents. *Fundam Appl Toxicol*, 37, 71.
 32. Oddi A, Mills CO, Custureri F, Di Nicola V, Elias E, Di Matteo G. (1996) Intraoperative biliary tree imaging with cholylyl-fluorescein: an experimental study in the rabbit. *Surg Laparosc Endosc*, 6, 198.
 33. Roman ID, Johnson GD, Coleman R. (1996) S-adenosyl-L-methionine prevents disruption of canalicular function and pericanalicular cytoskeleton integrity caused by cyclosporin A in isolated rat hepatocyte couplets. *Hepatology*, 24, 134.
 34. Stone V, Coleman R, Chipman JK. (1996) Comparison of the effects of redox cycling and arylating quinones on hepatobiliary function and glutathione homeostasis in rat hepatocyte couplets. *Toxicol Appl Pharmacol*, 138, 195.
 35. Baxter DJ, Rahman K, Bushell AJ, Mills CO, Elias E, Billington D. (1995) Biliary lipid output by isolated perfused rat livers in response to cholylyl-fluorescein. *Biochim Biophys Acta*, 1256, 374.
 36. Maglova LM, Jackson AM, Meng XJ, Carruth MW, Schteingart CD, Ton-Nu HT, Hofmann AF, Weinman SA. (1995) Transport characteristics of three fluorescent conjugated bile acid analogs in isolated rat hepatocytes and couplets. *Hepatology*, 22, 637.
 37. Roman ID, Coleman R. (1994) Disruption of canalicular function in isolated rat hepatocyte couplets caused by cyclosporin A. *Biochem Pharmacol*, 48, 2181.

38. Stone V, Johnson GD, Wilton JC, Coleman R, Chipman JK. (1994) Effect of oxidative stress and disruption of Ca²⁺ homeostasis on hepatocyte canalicular function in vitro. *Biochem Pharmacol*, 47, 625.
39. Weinman SA, Maglova LM. (1994) Free concentrations of intracellular fluorescent anions determined by cytoplasmic dialysis of isolated hepatocytes. *Am J Physiol*, 267, G922.
40. Wilton JC, Matthews GM, Burgoyne RD, Mills CO, Chipman JK, Coleman R. (1994) Fluorescent choleric and cholestatic bile salts take different paths across the hepatocyte: transcytosis of glycolithocholate leads to an extensive redistribution of annexin II. *J Cell Biol*, 127, 401.
41. Wilton JC, Chipman JK, Lawson CJ, Strain AJ, Coleman R. (1993) Periportal- and perivenous-enriched hepatocyte couplets: differences in canalicular activity and in response to oxidative stress. *Biochem J*, 292 (Pt 3), 773.
42. Wilton JC, Coleman R, Lankester DJ, Chipman JK. (1993) Stability and optimization of canalicular function in hepatocyte couplets. *Cell Biochem Funct*, 11, 179.
43. Mills CO, Rahman K, Coleman R, Elias E. (1991) Cholyl-l-lysylfluorescein: synthesis, biliary excretion in vivo and during single-pass perfusion of isolated perfused rat liver. *Biochim Biophys Acta*, 1115, 151.