References for Product 41

- Georgieva I, Trendafilova N, Aquino A, Lischka H. (2005) Excited state properties of 7hydroxy-4-methylcoumarin in the gas phase and in solution. A theoretical study. J Phys Chem A Mol Spectrosc Kinet Environ Gen Theory, 109, 11860.
- 2. Kumar S, Singh BK, Kalra N, Kumar V, Kumar A, Prasad AK, Raj HG, Parmar VS, Ghosh B. (2005) Novel thiocoumarins as inhibitors of TNF-alpha induced ICAM-1 expression on human umbilical vein endothelial cells (HUVECs) and microsomal lipid peroxidation. Bioorg Med Chem, 13, 1605.
- 3. Gleason NJ, Carbeck JD. (2004) Measurement of enzyme kinetics using microscale steady-state kinetic analysis. Langmuir, 20, 6374.
- 4. Kostova IP, Manolov, II, Radulova MK. (2004) Stability of the complexes of some lanthanides with coumarin derivatives. I. Cerium(III)-4-methyl-7-hydroxycoumarin. Acta Pharm, 54, 37.
- 5. Kawase M, Tanaka T, Sohara Y, Tani S, Sakagami H, Hauer H, Chatterjee SS. (2003) Structural requirements of hydroxylated coumarins for in vitro anti-Helicobacter pylori activity. In Vivo, 17, 509.
- 6. Kurosaki H, Sharma RK, Otsuka M, Goto M. (2003) Crystal structure of 7,8-dihydroxy-4-methylcoumarin. Anal Sci, 19, 647.
- 7. Yang J, Peng RX, Kong R, Yu JP. (2001) [Effects of 18 alpha-glycyrrhizic acid on rat liver cytochrome P450 isoenzymes and phase II transferase]. Yao Xue Xue Bao, 36, 321.
- 8. Everett SA, Swann E, Naylor MA, Stratford MR, Patel KB, Tian N, Newman RG, Vojnovic B, Moody CJ, Wardman P. (2002) Modifying rates of reductive elimination of leaving groups from indolequinone prodrugs: a key factor in controlling hypoxia-selective drug release. Biochem Pharmacol, 63, 1629.
- 9. Chilvers KF, Perry JD, James AL, Reed RH. (2001) Synthesis and evaluation of novel fluorogenic substrates for the detection of bacterial beta-galactosidase. J Appl Microbiol, 91. 1118.
- 10. Liu ZQ, Yu W, Liu ZL. (1999) Antioxidative and prooxidative effects of coumarin derivatives on free radical initiated and photosensitized peroxidation of human low-density lipoprotein. Chem Phys Lipids, 103, 125.
- 11. Ciotti M, Marrone A, Potter C, Owens IS. (1997) Genetic polymorphism in the human UGT1A6 (planar phenol) UDP-glucuronosyltransferase: pharmacological implications. Pharmacogenetics, 7, 485.
- 12. Duhaiman AS. (1996) Inhibition of zeta-crystallin by Coumarins: a structure-activity study. J Protein Chem, 15, 261.
- 13. Chang WS, Chiang HC. (1995) Structure-activity relationship of coumarins in xanthine oxidase inhibition. Anticancer Res, 15, 1969.
- 14. Irshaid Y, Abu-Khalaf M. (1992) Lack of effect of certain histamine H2-receptor blockers on the glucuronidation of 7-hydroxy-4-methylcoumarin by human liver microsomes. Pharmacol Toxicol, 71, 294.
- 15. Irshaid Y, Gharaybeh K. (1991) Glucuronidation of 7-hydroxy-4-methylcoumarin by microsomes from normal and diseased human livers. Pharmacol Toxicol, 68, 212.
- Irshaid YM, Gharaybeh KI, Ammari FF, Rawashdeh NM. (1990) Glucuronidation of 7hydroxy-4-methylcoumarin by human liver microsomes. Inhibition by certain drugs. Eur J Drug Metab Pharmacokinet, 15, 295.
- 17. Kuromizu K, Shimokawa Y, Abe O, Izumiya N. (1985) New fluorogenic substrate for esterase activity of alpha-chymotrypsin and related enzymes. Anal Biochem, 151, 534.
- 18. Powell SC, Scaro J, Wilson E, Shihabi ZK. (1983) Assay of urinary N-acetyl-beta-glucosaminidase in a centrifugal analyzer. Clin Chem, 29, 1717.
- 19. Baggett N, Marsden BJ. (1982) Reinvestigation of the synthesis of 4-methylcoumarin-7-yl 5-acetamido-3,5-dideoxy-alpha-D-glycero-D-galacto-2-nonulopyranosidonic acid, a fluorogenic substrate for neuraminidase. Carbohydr Res, 110, 11.
- 20. Curzon G, Kantamaneni BD. (1977) Fluorimetric determination of plasma unesterified fatty acid. Clin Chim Acta, 76, 289.