

## **References for Products 5000 to 5299**

1. Blanco AM, Rausell L, Aguado B, Perez-Alonso M, Artero R. (2009) A FRET-based assay for characterization of alternative splicing events using peptide nucleic acid fluorescence in situ hybridization. *Nucleic Acids Res*, 37, e116.
2. Ekici OD, Zhu J, Wah Chung IY, Paetzel M, Dalbey RE, Pei D. (2009) Profiling the substrate specificity of viral protease VP4 by a FRET-based peptide library approach. *Biochemistry*, 48, 5753.
3. Bruno JG, Carrillo MP, Phillips T. (2008) Development of DNA aptamers to a foot-and-mouth disease peptide for competitive FRET-based detection. *J Biomol Tech*, 19, 109.
4. Nettels D, Hoffmann A, Schuler B. (2008) Unfolded protein and peptide dynamics investigated with single-molecule FRET and correlation spectroscopy from picoseconds to seconds. *J Phys Chem B*, 112, 6137.
5. Oquare BY, Taylor JS. (2008) Synthesis of peptide nucleic acid FRET probes via an orthogonally protected building block for post-synthetic labeling of peptide nucleic acids at the 5-position of uracil. *Bioconjug Chem*, 19, 2196.
6. White BR, Liljestrand HM, Holcombe JA. (2008) A 'turn-on' FRET peptide sensor based on the mercury binding protein MerP. *Analyst*, 133, 65.
7. Alves FM, Hirata IY, Gouvea IE, Alves MF, Meldal M, Bromme D, Juliano L, Juliano MA. (2007) Controlled peptide solvation in portion-mixing libraries of FRET peptides: improved specificity determination for Dengue 2 virus NS2B-NS3 protease and human cathepsin S. *J Comb Chem*, 9, 627.
8. Deng W, Cao A, Lai L. (2007) Detecting the inter-peptide arrangement and maturation process of transthyretin (105-115) amyloid fibril using a FRET pair with short Forster distance. *Biochem Biophys Res Commun*, 362, 689.
9. Nakamura C, Miyamoto C, Obataya I, Takeda S, Yabuta M, Miyake J. (2007) Enzymatic nanolithography of FRET peptide layer using V8 protease-immobilized AFM probe. *Biosens Bioelectron*, 22, 2308.
10. Usui K, Takahashi M, Nokihara K, Mihara H. (2004) Peptide arrays with designed alpha-helical structures for characterization of proteins from FRET fingerprint patterns. *Mol Divers*, 8, 209.
11. Sun XG, Song G, Liu JH, An HY, Li HL, Xing FY, Zhang LH, Jiang Y. (2003) Construction of signal peptide-FRET fusion expression vectors and their expressions in NIH3T3 cells. *Di Yi Jun Yi Da Xue Xue Bao*, 23, 310.
12. Arslanbaeva LR, Zherdeva VV, Ivashina TV, Vinokurov LM, Rusanov AL, Savitskii AP. [Genetically encoded FRET-pair on the basis of terbium-binding peptide and red fluorescent protein]. *Prikl Biokhim Mikrobiol*, 46, 166.
13. Fu RH, Liu SP, Ou CW, Huang CM, Wang YC. Spatial control of cells, peptide delivery and dynamic monitoring of cellular physiology with chitosan-assisted dual color quantum dot FRET peptides. *Acta Biomater*, 6, 3621.
14. Lee H, Kim IK, Park TG. Intracellular trafficking and unpacking of siRNA/quantum dot-PEI complexes modified with and without cell penetrating peptide: confocal and flow cytometric FRET analysis. *Bioconjug Chem*, 21, 289.
15. Mathur N, Aneja A, Bhatnagar PK, Mathur PC. Detection of DNA Sequence with Enhanced Sensitivity and Higher FRET Efficiency Using a Light-Emitting Polymer, Peptide Nucleic Acid Probe and Anionic Surfactant System. *J Biomater Sci Polym Ed*.
16. Oliveira M, Torquato RJ, Alves MF, Juliano MA, Bromme D, Barros NM, Carmona AK. Improvement of cathepsin S detection using a designed FRET peptide based on putative natural substrates. *Peptides*, 31, 562.
17. Sakurai T, Iwasaki T, Okuno T, Kawata Y, Kise N. Evaluation of Abeta fibrillization inhibitory effect by a PEG-peptide conjugate based on an Abeta peptide fragment with intramolecular FRET. *Chem Commun (Camb)*, 47, 4709.